

# ANZSCP



## 15<sup>th</sup> ANNUAL MEETING

December 4 to 6  
1998

The University of Western Australia, Perth


---

# Programme & Abstracts


ISBN 0 86422 880 5

---

AUSTRALIA AND NEW ZEALAND SOCIETY FOR COMPARATIVE PHYSIOLOGY AND BIOCHEMISTRY  
proudly sponsored by: The University of Western Australia, Physiological Zoology/University of Chicago Press, Journal of Comparative  
Physiology/Springer, Royal Society of Western Australia: affiliated with IUBS and Comparative Biochemistry and Physiology




The 15<sup>th</sup> annual meeting of the Australia and New Zealand Society for Comparative Biochemistry and Physiology has benefited substantially by generous contributions from the following:



The University of Western Australia, Vice-Chancellory  
The University of Western Australia, Faculty of Science  
The Royal Society of Western Australia  
Springer Publishing (*Journal of Comparative Physiology*)  
University of Chicago Press (*Physiological and Biochemical Zoology*)  
Pergamon Press (*Comparative Biochemistry and Physiology*)


ANZSCP Proceedings,  
15<sup>th</sup> Annual Meeting, December 4-6, 1998  
The University of Western Australia, Nedlands WA 6907  
Organised by Philip C. Withers  
ISBN 0 86422 880 5

# PROGRAM



## FRIDAY December 4

|       |   |   |                                       |
|-------|---|---|---------------------------------------|
| 12:00 | <b>REGISTRATION and BARBEQUE</b>  |   | University House                      |
| 14:10 | <b>WELCOME</b>  |   | Zoology Lecture Theatre               |
|       | <b>SESSION I Metabolic Plasticity</b>   | <b>Chair: Mike Guppy</b>  | Zoology Lecture Theatre               |
| 14:20 | <u>Philip Withers</u>   | Overview of metabolic depression in ectotherms and endotherms   |                                       |
| 14:40 | <u>Fritz Geiser, G. Körtner &amp; I. Schmidt</u>                                | Effects of murine leptin on energetics and torpor of the dasyurid marsupial <i>Sminthopsis macroura</i>                           |                                       |
| 15:00 | <u>P. Hope, R. Baudinette, H. Turnbull, J. Kumaratilake, &amp; Gary Wittert</u> | Thermogenesis in marsupials: Uncoupling the evidence.   |                                       |
| 15:20 | <u>Graham Thompson</u>  | Comparing the metabolism of reptiles with varying body mass   |                                       |
| 15:40 | <b>AFTERNOON TEA</b>  |   | Zoology quadrangle                    |
|       | <b>SESSION 2 Metabolic Plasticity</b>   | <b>Chair: Fritz Geiser</b>  | Zoology Lecture Theatre               |
| 16:00 | <u>Philip Bethge &amp; S. Nicol</u>   | Energy requirements for different activities, including the cost of foraging, in the platypus ( <i>Ornithorhynchus anatinus</i> ) |                                       |
| 16:20 | <u>Julian Pakay, A. Hobbs, P. Withers &amp; M. Guppy</u>                        | The down-regulation of protein synthesis during metabolic depression in the land snail <i>Helix aspersa</i>                       |                                       |
| 16:40 | <u>Mark Chappell &amp; K. Hammond</u>   | Organ system correlates with variation in aerobic performance in red junglefowl   |                                       |
| 17:00 | <b>POSTERS — CHEESE &amp; WINE SOCIAL</b>                                       |   | Zoology 1 <sup>st</sup> floor balcony |
|       | <u>D. Adams, J. Cockrem &amp; M. Potter</u>                                     | Corticosterone response of captive northern brown kiwi ( <i>Apteryx australis mantelli</i> ) to an acute stressor.                |                                       |
|       | <u>P. Bethge, H. Otley, S. Munks, &amp; S. Nicol</u>                            | The use of dataloggers to determine behavioural activity in the platypus ( <i>Ornithorhynchus anatinus</i> )                      |                                       |
|       | <u>P. Celi, D. Blache, G. Martin, P. Vercoe, R. Dynes &amp; R. Tellam</u>       | Intracerebral infusion of recombinant leptin inhibits LH secretion in male sheep by decreasing feed intake                        |                                       |
|       | <u>L. Cutler &amp; A. Patak</u>   | Myofibrillar protein composition of regenerating and pristine claw closer muscle of the yabby, <i>Cherax albidus</i>              |                                       |
|       | <u>J. Elias &amp; J. Baldwin</u>  | Anaerobic metabolism in the intertidal elephant snail, <i>Scutus antipodes</i> (Fissurellidae)                                    |                                       |
|       | <u>B. Gartrell, S. Jones, R. Brereton &amp; L. Astheimer</u>                    | An investigation of the factors causing the population decline of the swift parrot ( <i>Lathamus discolor</i> )                   |                                       |
|       | <u>F. Geiser &amp; D. Coburn</u>  | Energy expenditure and water uptake in free-ranging and captive blossom-bats, <i>Syconycteris australis</i> (Megachiroptera)      |                                       |
|       | <u>L. Holm, I. Malecki &amp; G. Martin</u>                                      | Effect of temperature and pH on motility and velocity of emu spermatozoa  |                                       |
|       | <u>T. Lamey &amp; A. Patak</u>  | The effects of moulting on muscle fibre characteristics of the yabby, <i>Cherax albidus</i>                                       |                                       |
|       | <u>N. Leelapiyanart &amp; H. Taylor</u>   | Respiratory responses of intertidal crab eggs to changes in oxygen tension  |                                       |
|       | <u>O. Lopatko, P. Wood, J. Joss, A. Smits &amp; C. Daniels</u>                  | Acetylcholine controls surfactant secretion by Type II pneumocytes isolated from Australian lungfish and American bullfrogs       |                                       |
|       | <u>S. Orgeig, S. Johnston, O. Lopatko &amp; C. Daniels</u>                      | Development of the pulmonary surfactant system in a bird and reptile  |                                       |
|       | <u>J. Van Cleeff, M. Blackberry &amp; G. Martin</u>                             | Hyperlipaemic emu serum is clarified for radioimmunoassay by polyethyleneglycol (PEG) precipitation                               |                                       |
|       | <u>J. Wilde &amp; P. Greenaway</u>  | Diet and assimilation in the robber crab, <i>Birgus latro</i> (L.)  |                                       |



## SATURDAY December 5

| SESSION 3 |  | Chair: Jamie O'Shea   | Zoology Lecture Theatre |
|-----------|--|---|-------------------------|
| 09:00     | <u>John Donald</u> , B. Davis & K. Minerds                                       | The effect of total water deprivation on the natriuretic peptide system of the hopping mouse, <i>Notomys alexis</i>   |                         |
| 09:20     | K. Bell & <u>Sue Jones</u>   | Changes in plasma concentrations of corticosterone and testosterone in response to capture stress in the lizard <i>Egernia whitii</i>                         |                         |
| 09:40     | <u>W. Callahan</u> & <u>Tes Toop</u>   | Fishing for mechanisms: the natriuretic peptide system and the gills of the hagfish   |                         |
| 10:00     | <u>Julie Roberts</u>   | Aldosterone and renal function in avian species   |                         |
| 10:20     | <u>Dennyse Newbound</u> , J. O'Shea & P. Withers                                 | Osmoregulation and ionoregulation in the Port Jackson shark, <i>Heterodontus portusjacksoni</i>   |                         |
| 10:40     | MORNING TEA  |   | Zoology quadrangle      |
| SESSION 4 |  | Chair: Tess Toop  | Zoology Lecture Theatre |
| 11:00     | <u>Judith Van Cleeff</u> , D. Blache, I. Malecki, and G. B. Martin               | Gonadal hormones are proximate factors controlling seasonal fat metabolism in adult male emus   |                         |
| 11:20     | <u>Perdita Hope</u> , W. Breed, H. Turnbull, J. Morley, M. Horowitz & G. Wittert | Gonadal steroids and alterations in photoperiod affect metabolic efficiency in female <i>S. crassicaudata</i>   |                         |
| 11:40     | <u>Colleen Veitch</u> , C. Sernia & R. Gemmell                                   | Prostaglandin F <sub>2α</sub> Receptors in the brushtail possum, <i>Trichosurus vulpecula</i> and the short-nosed handicoot, <i>Isodon macrourus</i> at birth |                         |
| 12:00     | <u>Bronwyn McAllan</u> & J. Joss   | Effects of oral administration of melatonin on the reproductive cycle of a small marsupial, <i>Antechinus stuartii</i>  |                         |
| 12:20     | <u>Agung Riono</u> , D. Blache, A. Dharmarajan & G. Martin                       | Ingestion of extract of leaves of mangosteen ( <i>Garcinia mangostana</i> Linn) induces germ cell apoptosis in rat testis                                     |                         |
| 12:40     | LUNCH  |   | Zoology quadrangle      |
| SESSION 5 |  | Chair: Peter Greenaway  | Zoology Lecture Theatre |
| 14:00     | <u>Robert Gemmell</u> & B. Harmon  | The function of the vaginal caecae of the marsupial bandicoot, <i>Isodon macrourus</i>  |                         |
| 14:20     | <u>Michel Beal</u> & N. Scott  | Plasma concentrations of calcium, magnesium and phosphate during development of tammar pouch-young (PY)   |                         |
| 14:40     | <u>Ashley Edwards</u> & S M Jones  | Steroid biosynthesis in a viviparous reptile, <i>Tiliqua nigrolutea</i> (Scincidae)   |                         |
| 15:00     | <u>Peta Edwards</u> , B. Tuch & M. Beal  | Abundance of endocrine cells in the pancreatic islets of macropods  |                         |
| 15:20     | K. McClelland & <u>Ian Hume</u>  | Selective retention of digesta in the caecum of bandicoots  |                         |
| 15:40     | AFTERNOON TEA  |   | Zoology quadrangle      |
| SESSION 6 |  | Chair: Ian Hume   | Zoology Lecture Theatre |
| 16:00     | <u>Jamie O'Shea</u>  | The innervation of the heart of the fat-tailed dunnart, <i>Sminthopsis crassicaudata</i>  |                         |
| 16:20     | <u>Suzanne Munns</u> & B. Evans  | Hypercapnia induced depression of ventilation in the Gippsland water dragon   |                         |
| 16:40     | <u>Anton Thompson</u> & J. O'Shea  | A cholinergically mediated adrenergic response in the heart of the horned shark, <i>Heterodontus portusjacksoni</i>   |                         |
| 17:00     | <u>Alan Roberts</u>  | Ecological correlates of locomotor performance in geckos  |                         |
| 18:30     | Pre-dinner drinks  |   | University House        |
| 19:00     | 1998 ANNUAL MEETING DINNER   |   | University House        |

**SUNDAY December 6**

| <b>SESSION 3</b> |  | <b>Chair: Don Bradshaw</b>   | <b>Zoology Lecture Theatre</b> |
|------------------|--|--|--------------------------------|
| 09:00            | <u>Willie van Aardt</u>                                | The $MO_2$ and metabolic release of $CO_2$ from labeled glucose and lactate in the exercising ghost crab, <i>Ocypode ceratophthalmus</i> |                                |
| 09:20            | <u>Philip Withers, G. Thompson &amp; R. Seymour</u>    | Metabolic physiology of the marsupial mole ( <i>Notoryctes caurinus</i> )  |                                |
| 09:40            | <u>John Baldwin, A. Gupta &amp; X. Inglesias</u>       | Scaling of anaerobic energy metabolism during tail flipping behaviour in the freshwater crayfish, <i>Cherax destructor</i>               |                                |
| 10:00            | <u>Michael Guppy</u>                                   | A comparative study of fuel choices by cells   |                                |
| 10:20            | <u>Joseph Hoh, Y. Kim, L. Sieber &amp; C. Lucas</u>    | Macropodids express alpha cardiac myosin in their jaw-closing muscles  |                                |
| 10:40            | <b>MORNING TEA</b>                                     |  | <b>Zoology quadrangle</b>      |
| <b>SESSION 4</b> |  | <b>Chair: Phil Withers</b>   | <b>Zoology Lecture Theatre</b> |
| 11:00            | <u>Trevor Anderson &amp; S. Smith</u>                  | The relationship between histidase activity and trophic habit in surgeon fishes  |                                |
| 11:20            | <u>Don Bradshaw &amp; F. Bradshaw</u>                  | Measurement of nectar and pollen intake of free-ranging marsupial Honey possums ( <i>Tarsipes rostratus</i> )                            |                                |
| 11:40            | <u>Patrick Spanoghe, G. Davidson &amp; B. Paterson</u> | Effect of aerial exposure on the western rock lobster, <i>Panulirus cygnus</i>   |                                |
| 12:00            | <u>Stewart Nicol &amp; N. Andersen</u>                 | Using dataloggers to investigate hibernation in echidnas   |                                |
| 12:20            | <u>James Pearson</u>                                   | Physiological precocity in an altricial parrot   |                                |
| 12:40            | <b>Award of Student Prizes</b>                         |  | <b>Zoology Lecture Theatre</b> |
| 13:00            | <b>DISCUSSION SESSION - 1999 Meeting</b>               |  | <b>Zoology Lecture Theatre</b> |

**Australia  
New**

**Society for  
Comparative  
Physiology &  
Biochemistry**



### Information Office

Unipark Phone 9380 1229  
 9380 3018  
 Security Phone 9380 3020 - all hours  
 9380 2222 - emergencies only  
 1800 655 222 - free call from  
 all public phones to security



### CAR PARKS AND FACILITIES

Please read the signs before you park your vehicle\*

- Permit Parking** [shaded box]
- Yellow (students) 1, 4, 8, 9, 12, 14, 26, 31  
 Red (staff) 1, 3, 6-15, 18-21, 24-27, 29-32  
 Car pooling 4, 14 (car pooling permit required)  
 University House 2 (members and guests - permit required)  
 Disabled 1, 2, 3, 5, 8, 9, 11, 13 - 16, 18, 19, 30, 31 (UWA disabled or ACROD permit required)
- Reserved bays** reserved at all times  
**Service bays** 1-3, 5, 4, 8, 9, 11, 12, 13, 15, 16, 18, 19, 20, 22, 23, 30, 31 (place business/authority card on dashboard)
- Pay Parking** [shaded box]
- Short Term 2 hours - \$1.00 per hour - 4  
 Short Term 1 hour - \$1.00 per hour - 1, 3, 11, 20  
 All Day \*30 cents per hour - 9, 17, 21, 23, 31  
 (\* expected to rise to 50 cents in 1998)
- Motorcycle bays** 1, 3, 4, 8, 11, 13, 14, 15, 16, 18, 31  
 (no permit required)  
**Bus Parking** east of Music building - phone 9380 3020
- Bus Routes to Campus**  
 University service No 78, 79-74, 77, 102-4, 300-202, 308-9
- Disabled Toilet** [shaded box]   
**Public Toilet** [shaded box]   
**Public Telephones** [shaded box]   
**Bicycle Rack** [shaded box]   
**Bus Stop** [shaded box] 



**THE UNIVERSITY OF WESTERN AUSTRALIA**

## ABSTRACTS OF TALKS

### The relationship between histidase activity and trophic habit in surgeon fishes

Trevor A. Anderson and Shawn Smith

Department of Aquaculture, James Cook University, Townsville, QLD 4811  
Trevor.Anderson@jcu.edu.au

Herbivorous animals survive on a diet which is relatively poor in protein and essential amino acids (EAA) and may therefore possess mechanisms that protect the EAA from degradation by modifications in the  $K_m$  and  $V_{max}$  of the key degradative enzymes. We expected that the initial enzyme in the degradative pathway of an essential amino acid would have a lower  $K_m$  and a higher  $V_{max}$  in animals that receive an adequate supply of essential amino acids (carnivores). Our initial test of this hypothesis focussed on histidase, the primary enzyme in histidine degradation, as histidine levels are generally tightly regulated in fish. Individuals of five closely related species of tropical surgeon fish with a range of trophic habits were captured by spear, dissected and liver and muscle tissue stored in liquid nitrogen until assay. Homogenates of tissue were prepared and assayed for histidase using the method of Chiu *et al.* (1984).

The carnivorous *Naso hexacanthus* and *N. vlamingii* and the fermenting herbivore *N. unicornus* had lower  $K_m$  and  $V_{max}$  for liver histidase than the detritivorous *Ctenochaetus striatus* which had lower values than the non-fermenting herbivore *Acanthurus lineatus*. These data support the hypothesis with regard to  $K_m$  but not with regard to  $V_{max}$ .

Chiu, Y.N., Austic, R.E. & Rumsay, G.L. (1984) *Comp. Biochem. Physiol.* 78B, 777-783.

### Scaling of anaerobic energy metabolism during tail flipping behaviour in the freshwater crayfish, *Cherax destructor*

J. Baldwin, A. Gupta and X. Inglesias

Department of Biological Sciences, Monash University, Clayton, VIC 3168  
joha.baldwin@sci.monash.edu.au

Yabbies (*Cherax destructor*) display an escape behaviour in which rapid bursts of tail flipping propel the animal backwards. This is a classic example of anaerobic muscle work, initially powered by phosphagen hydrolysis then switching to anaerobic glycolysis once arginine phosphate reserves are depleted. We have investigated the effect of animal size on anaerobic scope (maximum rate of energy production) and anaerobic capacity (total energy used to reach exhaustion) during this escape behaviour.

The positive allometry of anaerobic scope, reflected in the activities of rate limiting glycolytic enzymes, scales with the increased power required by larger animals to overcome drag during locomotion through water. Exercise time and number of tail flips required to reach exhaustion showed positive allometry, as did anaerobic glycolytic capacity. However, the contribution of phosphagen hydrolysis to anaerobic capacity was independent of body mass. Limits to anaerobic capacity are not set by fuel stores, but may involve inhibition of glycolytic enzyme activity at the low pH values reached in tail muscle of exhausted animals. It is suggested that this positive allometry of anaerobic capacity observed during enforced exercise may not be utilised routinely in nature because of metabolic constraints imposed during recovery of large animals.

## Plasma concentrations of calcium, magnesium and phosphate during development of Tammar pouch-young (PY)

A. M. Beal and N. A. Scott

School of Biological Science, University of New South Wales, Sydney, NSW 2052  
m.beal@unsw.edu.au

At birth, marsupial neonates lack calcified bones. Whether they are born with some capacity to regulate calcium and phosphorus or when this develops during pouch-life is unknown. We have measured plasma and milk concentrations of Ca, Mg, PO<sub>4</sub> and other ions from mothers and PY starting at 25 d post-birth. PY plasma [Ca] increased progressively from 25 to 150 d post-natum but remained lower than maternal plasma Ca ( $P < 0.001$ ) up to 100 d. In contrast, PY plasma was hypermagnesian to maternal plasma being near constant up to 100 d post-birth but rising between 100 and 200 d. Despite milk having substantially higher levels of both elements than either maternal or PY plasma, the concentrations of Ca and Mg in milk were not correlated with plasma levels of mothers ( $P = 0.4$  and  $0.93$ ) or PY ( $P = 0.18$  and  $0.54$ ). PY were hyperphosphataemic relative to the mothers throughout pouch-life with PY plasma concentrations increasing over the 100-250 d post-birth. Milk [PO<sub>4</sub>] were much higher than that in plasma from the mothers or PY and was negatively correlated with maternal plasma ( $P = 0.023$ ) and positively correlated with PY plasma ( $P = 0.005$ ) throughout pouch life. The data suggest that PY gain control of Ca homeostasis earlier than that of Mg or PO<sub>4</sub>.

## Changes in plasma concentrations of corticosterone and testosterone in response to capture stress in the lizard *Egernia whitii*

K. Bell and S. M. Jones

School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001  
S.M.Jones@utas.edu.au

In vertebrates, exposure to a stressor generally stimulates the secretion of corticosteroids by the adrenal cortex. Often, an increase in plasma corticosteroids is followed by a decrease in plasma sex steroid concentrations; however, in reptiles, these generalisations do not apply to all species. Males of the lizard *Egernia whitii* were captured in the field and bled at 0, 10, 60 or 240 minutes after capture, each animal being sampled once only. The response to acute capture stress was reflected a rapid increase in plasma corticosterone concentrations, with a significant variation after ten minutes, and concentrations remaining high at 240 min. Plasma testosterone concentrations also changed dramatically, but at a slower rate. Gonadal interactions in *E. whitii* were also investigated during long term captivity, and through the reproductive cycle. There were no significant differences between males and females in plasma concentrations of corticosterone through the season, with lowest values occurring during the postpartum period. These results were unexpected; the lack of sex-related differences in plasma corticosteroid concentrations may reflect the social nature of this species.



**Energy requirements for different activities, including  
the cost of foraging, in the platypus  
(*Ornithorhynchus anatinus*)**

Phillip Bethge and Stewart Nicol

*Division of Anatomy and Physiology, University of Tasmania, GPO Box 252-24,  
Hobart, TAS 7001  
p\_bethge@utas.edu.au*

We measured the energy requirements of freely-swimming platypuses in a 11 m<sup>3</sup> swim tank using respirometry. In the tank animals foraged voluntarily while all activities and gaseous exchange were closely monitored. In addition we used a conventional treadmill to determine energy requirements for walking. Energy requirements of the active platypus were found to be higher than maximal metabolic rates for the animal reported in the past. Energy requirements for foraging averaged 8.6 W/kg (n=6) for dives lasting the normal dive duration of 20 to 40 s observed in the wild and increased slightly with decreasing dive duration. The metabolic rate for walking was found to be 8.7 W/kg (speed 0.2 m/s, n=8). Resting rates of the platypus were found to increase with decreasing air and water temperature respectively. Minimal energy requirements for resting under the water surface (wedging) amounted to 3.6 W/kg (water temp 14°C). Minimal metabolic rate of platypuses resting on the water surface was found to be 3.9 W/kg (water temp 20°C) while minimal resting rate on land was 2.1 W/kg (air temp 26°C).

**Measurement of nectar and pollen intake  
of free-ranging marsupial honey possums  
(*Tarsipes rostratus*)**

S. D. Bradshaw and F. J. Bradshaw

*Department of Zoology, The University of Western Australia, Perth, WA 6907  
Don.Bradshaw@uwa.edu.au*

Turnover rates of a combination of stable and radioactive isotopes (tritium, oxygen-18 and sodium-22) have been used to estimate daily intake of pollen and nectar in free-ranging honey possums in Scott National Park in the extreme southwest of Western Australia. The Field Metabolic Rate (FMR) is measured using doubly-labelled water and nectar intake is estimated independently from the turnover of both water and sodium. Nectar energy intake is then estimated from the sugar concentration of fresh nectar collected from the inflorescences on which the honey possums have been feeding. Pollen intake is finally estimated as the difference between the energy derived from nectar and the FMR. The method assumes that honey possums normally do not drink free-water in the field, that all the sodium in their diet is derived from nectar, and that individuals are in energetic balance over the measurement period. These assumptions have been tested and found to be reasonably robust, except during periods of heavy rain when some free-water intake may occur. Data from some 30 individuals show that nectar intake is reasonably constant at 6-7 ml.day<sup>-1</sup> for individuals with an average mass of 9 g. Daily pollen intake varies considerably between individuals, however, averaging 510 mg.day<sup>-1</sup> but ranging from 0 to as much as 2 g.day<sup>-1</sup>. Laboratory studies with a captive colony of honey possums show that the minimum nitrogen requirement for the maintenance of balance in this species is approximately 115 mg.kg<sup>-0.75</sup>.day<sup>-1</sup>, well below that recorded for macropodid marsupials but very similar to that of the sugar glider, *Petaurus breviceps*, which also feeds on a high carbohydrate diet.

**Fishing for mechanisms:  
the natriuretic peptide system  
and the gills of the hagfish**

**Will Callahan and Tes Toop**

*School of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217  
ttop@deakin.edu.au*

Previous studies have identified a natriuretic peptide (NP) system in the hagfish. This earlier research focussed on the Atlantic hagfish, *Myxine glutinosa*. Recently, whole gill perfusion studies and myography of afferent and efferent branchial arteries have implicated NPs in the control of blood flow through the gills of the New Zealand hagfish, *Eptatretus cirrhatus* (Forster, Glover and Simpson, XIII Int. Congress Comp. Endocrinol. 1997). However, the NP receptors that would mediate this regulation have not been identified in this species. The current study demonstrates the presence of NP binding sites in the gills of *E. cirrhatus* and examines whether NP actions in the gills of this hagfish could be mediated by guanylate cyclase receptors, similar to those found in jawed vertebrates.

**Organ system correlates with variation  
in aerobic performance in red junglefowl**

**M. A. Chappell and K. Hammond**

*Department of Biology, University of California, Riverside, CA (USA)  
chappell@citrus.ucr.edu*

Limits to whole-animal performance may be set by peripheral effectors, central organs, or by all components in synchrony ("symmorphosis"). We examined these concepts using aerobic performance in red junglefowl as a model system. We measured individual variation in the masses of peripheral (pectoralis and leg muscles) and central organs (heart, lungs, gonads, liver, spleen, and gut), and searched for correlations with minimal and maximal aerobic metabolism (BMR and  $VO_{2max}$ ). We found substantial variance within and between sexes, but few consistent relationships between performance and organ size. On average, males had lower BMR and higher  $VO_{2max}$  than females, and had smaller visceral organs (gut, liver) and larger cardiovascular organs (heart, lung, haematocrit) and muscles. BMR was not correlated with any single organ mass in either sex (there was no relationship between female reproductive status and either BMR or  $VO_{2max}$ , although gonadal and oviduct mass varied more than 12-fold). In males only, BMR was correlated to combined gut mass. Male  $VO_{2max}$  residuals were correlated to heart and muscle mass; female  $VO_{2max}$  residuals were correlated to haematocrit. These results suggest that attaining high aerobic performance requires increased investment in muscles and in organs involved in oxygen delivery, but not in digestive organs.

**The effect of total water deprivation  
on the natriuretic peptide system of the hopping mouse,  
*Notomys alexis***

J. A. Donald, B. Davis and K. Minerds

School of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217  
jdonald@deakin.edu.au

The natriuretic peptide system counters hypervolaemia and concomitant hypertension by renal diuresis and natriuresis, and vasodilation. However, the importance of hypervolaemic regulation in mammals has been questioned, and in particular, the role of natriuretic peptides in the maintenance of normal fluid balance. We are using the hopping mouse, *Notomys alexis*, to investigate the function of natriuretic peptides in a desert-adapted mammal that can survive without free access to water. Atrial natriuretic peptide (ANP) cDNA was cloned from *N. alexis*, and the sequence of ANP (1-28) was identical to rat ANP. This information enabled a radioimmunoassay for rat ANP to be used to quantify ANP levels in the heart and plasma of *N. alexis*. *N. alexis*, of either sex, were divided into an experimental group (n=12) subject to seventeen days of total water deprivation (TWD), and a control group (n=12) that had free access to water. No significant difference in haematocrit, skeletal muscle and liver water content, and cardiac ANP levels was observed between control and experimental groups. However, plasma ANP and the density of ANP binding on renal glomeruli were significantly lower in TWD animals. This data suggests that the ANP system is down-regulated during TWD, probably to conserve salt and water.

**Steroid biosynthesis in a viviparous reptile,  
*Tiliqua nigrolutea* (Scincidae)**

A. Edwards, S. M. Jones and N. Davies

School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001  
Ashley.Edwards@utas.edu.au

Predominance of either the delta-4 or delta-5 pathway for gonadal steroid biosynthesis varies between vertebrate classes. More recent information suggests that variation between both sex and reproductive condition can also occur. We examined *in vitro* biosynthesis of progesterone (P4), androstendione (AD), dehydroepiandrosterone (DHA), testosterone (T) and 17 $\beta$ -oestradiol (E2) from pregnenolone (P5). Late pre-ovulatory (Oct) and post-partum (Mar) ovaries and final-spermatogenesis (Sep) and quiescent (Feb) testes of a viviparous skink, *Tiliqua nigrolutea* were used. Steroids were identified using various thin layer chromatography (TLC) solvent systems and HPLC (attached to a radiometric detector) elution times. A strong preference for the delta-4 pathway was apparent in both sexes and at both reproductive and quiescent times of year. However, for several key steroids, the relative proportions synthesised during incubation differed between reproductive conditions. Interestingly, we have demonstrated that ovarian follicles do not produce E2 from P5 *in vitro* at either time of year. The end product of P5 metabolism is instead another, more polar steroid, which is currently unidentified.

## Abundance of endocrine cells in the pancreatic islets of macropods

P. Edwards, B. E. Tuch and A. M. Beal

School of Biological Science, University of New South Wales and Dept of Endocrinology, Prince of Wales Hospital, Sydney, NSW 2052  
m.beal@unsw.edu.au

For more than twenty years, it has been believed that the islets of Langerhans from the macropod pancreas were essentially similar to those from the sheep pancreas in having a preponderance of  $\alpha$  cells (glucagon-secreting) and few  $\beta$  cells (insulin-secreting) in comparison with islets from monogastric animals. This was argued to be an adaptation to foregut fermentation. On finding that the islets of one tammur wallaby had high  $\beta$  and low  $\alpha$  cell numbers we have reassessed the proportions of the 4 cell types in islets from red and eastern grey kangaroos, tammars and sheep using immunohistological techniques. The results confirm that sheep islets have high  $\alpha$  cell (53.9  $\pm$  3.20 %) and low  $\beta$  cell numbers (29.1  $\pm$  1.61 %) with the remainder being somatostatin-secreting  $\delta$  cells (6.2  $\pm$  1.96 %) and pancreatic polypeptide secreting  $\phi$  cells (10.8  $\pm$  4.04 %). In contrast, reds, greys and tammars were similar in having nearly twice as many  $\beta$  cells as  $\alpha$  cells averaging 52.4  $\pm$  4.36 %  $\beta$ -cells, 28.9  $\pm$  2.27 %  $\alpha$ -cells, 9.2  $\pm$  1.33 %  $\delta$ -cells and 10.5  $\pm$  2.58 %  $\phi$ -cells. Apparently, high  $\alpha$  cell numbers are not an essential adjunct to being a foregut.

## Effects of murine leptin on energetics and torpor of the dasyurid marsupial *Sminthopsis macroura*

Fritz Geiser<sup>1</sup>, Gerhard Körtner<sup>1</sup> and Ingrid Schmidt<sup>2</sup>

<sup>1</sup>Zoology, School of Biological Sciences, University of New England, Armidale, NSW 2351; <sup>2</sup>Max-Planck-Institut für physiologische und klinische Forschung, W.G. Kerckhoff-Institut, D-61231, Bad Nauheim, Germany  
fgeiser@metz.une.edu.au

Leptin plays an important role in regulating body fat stores of placental mammals, but the contribution of changes in energy uptake and expenditure to this adjustment remains controversial. We were interested in how recombinant murine leptin would affect metabolic rate and body temperature of a marsupial mammal (*Sminthopsis macroura*, 25 g) known to display daily torpor. In a group of 8 animals, food-deprived for 1 day at 18 °C, leptin treatment (5  $\mu$ g/g body mass in Tris buffer) halved the duration of torpor bouts, and raised the average daily minimum body temperature by 4.5 °C and the minimum metabolic rate by 2.2-fold in comparison to control treatment with Tris buffer. Leptin treatment thus increased daily energy expenditure by 9% although during the activity phase metabolic rates and body temperatures were not raised. Body mass was also not affected. These findings suggest that leptin affects the adjustment of thermoregulatory energy expenditure during the rest phase. They show that the hormone has strong physiological effects despite the lack of thermogenetically active brown fat in marsupials and some structural differences between murine and marsupial leptin.