


ANZSCP



15th ANNUAL MEETING

December 4 to 6
1998

The University of Western Australia, Perth


Programme & Abstracts

ISBN 0 86422 880 5

AUSTRALIA AND NEW ZEALAND SOCIETY FOR COMPARATIVE PHYSIOLOGY AND BIOCHEMISTRY
proudly sponsored by: The University of Western Australia, Physiological Zoology/University of Chicago Press, Journal of Comparative Physiology/Springer, Royal Society of Western Australia: affiliated with IUBS and Comparative Biochemistry and Physiology




The 15th annual meeting of the Australia and New Zealand Society for Comparative Biochemistry and Physiology has benefited substantially by generous contributions from the following:



The University of Western Australia, Vice-Chancellory
The University of Western Australia, Faculty of Science
The Royal Society of Western Australia
Springer Publishing (*Journal of Comparative Physiology*)
University of Chicago Press (*Physiological and Biochemical Zoology*)
Pergamon Press (*Comparative Biochemistry and Physiology*)


ANZSCP Proceedings,
15th Annual Meeting, December 4-6, 1998
The University of Western Australia, Nedlands WA 6907
Organised by Philip C. Withers
ISBN 0 86422 880 5

PROGRAM



FRIDAY December 4

12:00	REGISTRATION and BARBEQUE	University House
14:10	WELCOME	Zoology Lecture Theatre
	SESSION I Metabolic Plasticity	Chair: Mike Guppy
		Zoology Lecture Theatre
14:20	<u>Philip Withers</u>	Overview of metabolic depression in ectotherms and endotherms
14:40	<u>Fritz Geiser, G. Körtner & I. Schmidt</u>	Effects of murine leptin on energetics and torpor of the dasyurid marsupial <i>Sminthopsis macroura</i>
15:00	<u>P. Hope, R. Baudinette, H. Turnbull, J. Kumaratilake, & Gary Wittert</u>	Thermogenesis in marsupials: Uncoupling the evidence.
15:20	<u>Graham Thompson</u>	Comparing the metabolism of reptiles with varying body mass
15:40	AFTERNOON TEA	Zoology quadrangle
	SESSION 2 Metabolic Plasticity	Chair: Fritz Geiser
		Zoology Lecture Theatre
16:00	<u>Philip Bethge & S. Nicol</u>	Energy requirements for different activities, including the cost of foraging, in the platypus (<i>Ornithorhynchus anatinus</i>)
16:20	<u>Julian Pakay, A. Hobbs, P. Withers & M. Guppy</u>	The down-regulation of protein synthesis during metabolic depression in the land snail <i>Helix aspersa</i>
16:40	<u>Mark Chappell & K. Hammond</u>	Organ system correlates with variation in aerobic performance in red junglefowl
17:00	POSTERS — CHEESE & WINE SOCIAL	Zoology 1st floor balcony
	<u>D. Adams, J. Cockrem & M. Potter</u>	Corticosterone response of captive northern brown kiwi (<i>Apteryx australis mantelli</i>) to an acute stressor.
	<u>P. Bethge, H. Otley, S. Munks, & S. Nicol</u>	The use of dataloggers to determine behavioural activity in the platypus (<i>Ornithorhynchus anatinus</i>)
	<u>P. Celi, D. Blache, G. Martin, P. Vercoe, R. Dynes & R. Tellam</u>	Intracerebral infusion of recombinant leptin inhibits LH secretion in male sheep by decreasing feed intake
	<u>L. Cutler & A. Patak</u>	Myofibrillar protein composition of regenerating and pristine claw closer muscle of the yabby, <i>Cherax albidus</i>
	<u>J. Elias & J. Baldwin</u>	Anaerobic metabolism in the intertidal elephant snail, <i>Scutus antipodes</i> (Fissurellidae)
	<u>B. Gartrell, S. Jones, R. Brereton & L. Astheimer</u>	An investigation of the factors causing the population decline of the swift parrot (<i>Lathamus discolor</i>)
	<u>F. Geiser & D. Coburn</u>	Energy expenditure and water uptake in free-ranging and captive blossom-bats, <i>Syconycteris australis</i> (Megachiroptera)
	<u>L. Holm, I. Malecki & G. Martin</u>	Effect of temperature and pH on motility and velocity of emu spermatozoa
	<u>T. Lamey & A. Patak</u>	The effects of moulting on muscle fibre characteristics of the yabby, <i>Cherax albidus</i>
	<u>N. Leelapiyanart & H. Taylor</u>	Respiratory responses of intertidal crab eggs to changes in oxygen tension
	<u>O. Lopatko, P. Wood, J. Joss, A. Smits & C. Daniels</u>	Acetylcholine controls surfactant secretion by Type II pneumocytes isolated from Australian lungfish and American bullfrogs
	<u>S. Orgeig, S. Johnston, O. Lopatko & C. Daniels</u>	Development of the pulmonary surfactant system in a bird and reptile
	<u>J. Van Cleeff, M. Blackberry & G. Martin</u>	Hyperlipaemic emu serum is clarified for radioimmunoassay by polyethyleneglycol (PEG) precipitation
	<u>J. Wilde & P. Greenaway</u>	Diet and assimilation in the robber crab, <i>Birgus latro</i> (L.)



SATURDAY December 5

SESSION 3		Chair: Jamie O'Shea	Zoology Lecture Theatre
09:00	<u>John Donald</u> , B. Davis & K. Minerds	The effect of total water deprivation on the natriuretic peptide system of the hopping mouse, <i>Notomys alexis</i>	
09:20	K. Bell & <u>Sue Jones</u>	Changes in plasma concentrations of corticosterone and testosterone in response to capture stress in the lizard <i>Egernia whitii</i>	
09:40	<u>W. Callahan</u> & <u>Tes Toop</u>	Fishing for mechanisms: the natriuretic peptide system and the gills of the hagfish	
10:00	<u>Julie Roberts</u>	Aldosterone and renal function in avian species	
10:20	<u>Dennyse Newbound</u> , J. O'Shea & P. Withers	Osmoregulation and ionoregulation in the Port Jackson shark, <i>Heterodontus portusjacksoni</i>	
10:40	MORNING TEA		Zoology quadrangle
SESSION 4		Chair: Tess Toop	Zoology Lecture Theatre
11:00	<u>Judith Van Cleeff</u> , D. Blache, I. Malecki, and G. B. Martin	Gonadal hormones are proximate factors controlling seasonal fat metabolism in adult male emus	
11:20	<u>Perdita Hope</u> , W. Breed, H. Turnbull, J. Morley, M. Horowitz & G. Wittert	Gonadal steroids and alterations in photoperiod affect metabolic efficiency in female <i>S. crassicaudata</i>	
11:40	<u>Colleen Veitch</u> , C. Sernia & R. Gemmell	Prostaglandin F _{2α} Receptors in the brushtail possum, <i>Trichosurus vulpecula</i> and the short-nosed handicoot, <i>Isodon macrourus</i> at birth	
12:00	<u>Bronwyn McAllan</u> & J. Joss	Effects of oral administration of melatonin on the reproductive cycle of a small marsupial, <i>Antechinus stuartii</i>	
12:20	<u>Agung Riono</u> , D. Blache, A. Dharmarajan & G. Martin	Ingestion of extract of leaves of mangosteen (<i>Garcinia mangostana</i> Linn) induces germ cell apoptosis in rat testis	
12:40	LUNCH		Zoology quadrangle
SESSION 5		Chair: Peter Greenaway	Zoology Lecture Theatre
14:00	<u>Robert Gemmell</u> & B. Harmon	The function of the vaginal caecae of the marsupial bandicoot, <i>Isodon macrourus</i>	
14:20	<u>Michel Beal</u> & N. Scott	Plasma concentrations of calcium, magnesium and phosphate during development of tammar pouch-young (PY)	
14:40	<u>Ashley Edwards</u> & S M Jones	Steroid biosynthesis in a viviparous reptile, <i>Tiliqua nigrolutea</i> (Scincidae)	
15:00	<u>Peta Edwards</u> , B. Tuch & M. Beal	Abundance of endocrine cells in the pancreatic islets of macropods	
15:20	K. McClelland & <u>Ian Hume</u>	Selective retention of digesta in the caecum of bandicoots	
15:40	AFTERNOON TEA		Zoology quadrangle
SESSION 6		Chair: Ian Hume	Zoology Lecture Theatre
16:00	<u>Jamie O'Shea</u>	The innervation of the heart of the fat-tailed dunnart, <i>Sminthopsis crassicaudata</i>	
16:20	<u>Suzanne Munns</u> & B. Evans	Hypercapnia induced depression of ventilation in the Gippsland water dragon	
16:40	<u>Anton Thompson</u> & J. O'Shea	A cholinergically mediated adrenergic response in the heart of the horned shark, <i>Heterodontus portusjacksoni</i>	
17:00	<u>Alan Roberts</u>	Ecological correlates of locomotor performance in geckos	
18:30	Pre-dinner drinks		University House
19:00	1998 ANNUAL MEETING DINNER		University House

SUNDAY December 6

SESSION 3		Chair: Don Bradshaw	Zoology Lecture Theatre
09:00	<u>Willie van Aardt</u>	The MO_2 and metabolic release of CO_2 from labeled glucose and lactate in the exercising ghost crab, <i>Ocypode ceratophthalmus</i>	
09:20	<u>Philip Withers, G. Thompson & R. Seymour</u>	Metabolic physiology of the marsupial mole (<i>Notoryctes caurinus</i>)	
09:40	<u>John Baldwin, A. Gupta & X. Inglesias</u>	Scaling of anaerobic energy metabolism during tail flipping behaviour in the freshwater crayfish, <i>Cherax destructor</i>	
10:00	<u>Michael Guppy</u>	A comparative study of fuel choices by cells	
10:20	<u>Joseph Hoh, Y. Kim, L. Sieber & C. Lucas</u>	Macropodids express alpha cardiac myosin in their jaw-closing muscles	
10:40	MORNING TEA		Zoology quadrangle
SESSION 4		Chair: Phil Withers	Zoology Lecture Theatre
11:00	<u>Trevor Anderson & S. Smith</u>	The relationship between histidase activity and trophic habit in surgeon fishes	
11:20	<u>Don Bradshaw & F. Bradshaw</u>	Measurement of nectar and pollen intake of free-ranging marsupial Honey possums (<i>Tarsipes rostratus</i>)	
11:40	<u>Patrick Spanoghe, G. Davidson & B. Paterson</u>	Effect of aerial exposure on the western rock lobster, <i>Panulirus cygnus</i>	
12:00	<u>Stewart Nicol & N. Andersen</u>	Using dataloggers to investigate hibernation in echidnas	
12:20	<u>James Pearson</u>	Physiological precocity in an altricial parrot	
12:40	Award of Student Prizes		Zoology Lecture Theatre
13:00	DISCUSSION SESSION - 1999 Meeting		Zoology Lecture Theatre

Australia
New

Society for
Comparative
Physiology &
Biochemistry



Information Office
 Unipark Phone 9380 1229
 9380 3018
 Security Phone 9380 3020 - all hours
 9380 2222 - emergencies only
 1800 655 222 - free call from
 all public phones to security



CAR PARKS AND FACILITIES
 Please read the signs before you park your vehicle*

- Permit Parking**
- Yellow (students) 1, 4, 8, 9, 12, 14, 26, 31
 - Red (staff) 1, 3, 6-15, 18-21, 24-27, 29-32
 - Car pooling 4, 14 (car pooling permit required)
 - University House 2 (members and guests - permit required)
 - Disabled 1, 2, 3, 5, 8, 9, 11, 13, 16, 18, 19, 30, 31 (UWA disabled or ACROD permit required)
- Reserved bays** reserved at all times
- Service bays** 1-3, 5, 4, 8, 9, 11, 12, 13, 15, 16, 18, 19, 20, 22, 27, 30, 31 (place business/authority card on dashboard)
- Pay Parking**
- Short Term 2 hours - \$1.00 per hour - 4
 - Short Term 1 hour - \$1.00 per hour - 1, 3, 11, 20
 - All Day *30 cents per hour - 9, 17, 21, 23, 31 (* expected to rise to 50 cents in 1998)
- Motorcycle bays** 1, 3, 4, 8, 11, 13, 14, 15, 16, 18, 31
 (no permit required)
- Bus Parking** east of Music building - phone 9380 3020
- Bus Routes to Campus**
 University service No 78, 79-74, 77, 102-4, 300-202, 308-9
- Disabled Toilet**
- Public Toilet**
- Public Telephones**
- Bicycle Rack**
- Bus Stop**



THE UNIVERSITY OF WESTERN AUSTRALIA

ABSTRACTS OF TALKS

The relationship between histidase activity and trophic habit in surgeon fishes

Trevor A. Anderson and Shawn Smith

Department of Aquaculture, James Cook University, Townsville, QLD 4811
Trevor.Anderson@jcu.edu.au

Herbivorous animals survive on a diet which is relatively poor in protein and essential amino acids (EAA) and may therefore possess mechanisms that protect the EAA from degradation by modifications in the K_m and V_{max} of the key degradative enzymes. We expected that the initial enzyme in the degradative pathway of an essential amino acid would have a lower K_m and a higher V_{max} in animals that receive an adequate supply of essential amino acids (carnivores). Our initial test of this hypothesis focussed on histidase, the primary enzyme in histidine degradation, as histidine levels are generally tightly regulated in fish. Individuals of five closely related species of tropical surgeon fish with a range of trophic habits were captured by spear, dissected and liver and muscle tissue stored in liquid nitrogen until assay. Homogenates of tissue were prepared and assayed for histidase using the method of Chiu *et al.* (1984).

The carnivorous *Naso hexacanthus* and *N. vlamingii* and the fermenting herbivore *N. unicornus* had lower K_m and V_{max} for liver histidase than the detritivorous *Ctenochaetus striatus* which had lower values than the non-fermenting herbivore *Acanthurus lineatus*. These data support the hypothesis with regard to K_m but not with regard to V_{max} .

Chiu, Y.N., Austic, R.E. & Rumsay, G.L. (1984) *Comp. Biochem. Physiol.* 78B, 777-783.

Scaling of anaerobic energy metabolism during tail flipping behaviour in the freshwater crayfish, *Cherax destructor*

J. Baldwin, A. Gupta and X. Inglesias

Department of Biological Sciences, Monash University, Clayton, VIC 3168
joha.baldwin@sci.monash.edu.au

Yabbies (*Cherax destructor*) display an escape behaviour in which rapid bursts of tail flipping propel the animal backwards. This is a classic example of anaerobic muscle work, initially powered by phosphagen hydrolysis then switching to anaerobic glycolysis once arginine phosphate reserves are depleted. We have investigated the effect of animal size on anaerobic scope (maximum rate of energy production) and anaerobic capacity (total energy used to reach exhaustion) during this escape behaviour.

The positive allometry of anaerobic scope, reflected in the activities of rate limiting glycolytic enzymes, scales with the increased power required by larger animals to overcome drag during locomotion through water. Exercise time and number of tail flips required to reach exhaustion showed positive allometry, as did anaerobic glycolytic capacity. However, the contribution of phosphagen hydrolysis to anaerobic capacity was independent of body mass. Limits to anaerobic capacity are not set by fuel stores, but may involve inhibition of glycolytic enzyme activity at the low pH values reached in tail muscle of exhausted animals. It is suggested that this positive allometry of anaerobic capacity observed during enforced exercise may not be utilised routinely in nature because of metabolic constraints imposed during recovery of large animals.

Plasma concentrations of calcium, magnesium and phosphate during development of Tammar pouch-young (PY)

A. M. Beal and N. A. Scott

School of Biological Science, University of New South Wales, Sydney, NSW 2052
m.beal@unsw.edu.au

At birth, marsupial neonates lack calcified bones. Whether they are born with some capacity to regulate calcium and phosphorus or when this develops during pouch-life is unknown. We have measured plasma and milk concentrations of Ca, Mg, PO₄ and other ions from mothers and PY starting at 25 d post-birth. PY plasma [Ca] increased progressively from 25 to 150 d post-natum but remained lower than maternal plasma Ca ($P < 0.001$) up to 100 d. In contrast, PY plasma was hypermagnesian to maternal plasma being near constant up to 100 d post-birth but rising between 100 and 200 d. Despite milk having substantially higher levels of both elements than either maternal or PY plasma, the concentrations of Ca and Mg in milk were not correlated with plasma levels of mothers ($P = 0.4$ and 0.93) or PY ($P = 0.18$ and 0.54). PY were hyperphosphataemic relative to the mothers throughout pouch-life with PY plasma concentrations increasing over the 100-250 d post-birth. Milk [PO₄] were much higher than that in plasma from the mothers or PY and was negatively correlated with maternal plasma ($P = 0.023$) and positively correlated with PY plasma ($P = 0.005$) throughout pouch life. The data suggest that PY gain control of Ca homeostasis earlier than that of Mg or PO₄.

Changes in plasma concentrations of corticosterone and testosterone in response to capture stress in the lizard *Egernia whitii*

K. Bell and S. M. Jones

School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001
S.M.Jones@utas.edu.au

In vertebrates, exposure to a stressor generally stimulates the secretion of corticosteroids by the adrenal cortex. Often, an increase in plasma corticosteroids is followed by a decrease in plasma sex steroid concentrations; however, in reptiles, these generalisations do not apply to all species. Males of the lizard *Egernia whitii* were captured in the field and bled at 0, 10, 60 or 240 minutes after capture, each animal being sampled once only. The response to acute capture stress was reflected a rapid increase in plasma corticosterone concentrations, with a significant variation after ten minutes, and concentrations remaining high at 240 min. Plasma testosterone concentrations also changed dramatically, but at a slower rate. Gonadal interactions in *E. whitii* were also investigated during long term captivity, and through the reproductive cycle. There were no significant differences between males and females in plasma concentrations of corticosterone through the season, with lowest values occurring during the postpartum period. These results were unexpected; the lack of sex-related differences in plasma corticosteroid concentrations may reflect the social nature of this species.

**Energy requirements for different activities, including
the cost of foraging, in the platypus
(*Ornithorhynchus anatinus*)**

Phillip Bethge and Stewart Nicol

*Division of Anatomy and Physiology, University of Tasmania, GPO Box 252-24,
Hobart, TAS 7001
p_bethge@utas.edu.au*

We measured the energy requirements of freely-swimming platypuses in a 11 m³ swim tank using respirometry. In the tank animals foraged voluntarily while all activities and gaseous exchange were closely monitored. In addition we used a conventional treadmill to determine energy requirements for walking. Energy requirements of the active platypus were found to be higher than maximal metabolic rates for the animal reported in the past. Energy requirements for foraging averaged 8.6 W/kg (n=6) for dives lasting the normal dive duration of 20 to 40 s observed in the wild and increased slightly with decreasing dive duration. The metabolic rate for walking was found to be 8.7 W/kg (speed 0.2 m/s, n=8). Resting rates of the platypus were found to increase with decreasing air and water temperature respectively. Minimal energy requirements for resting under the water surface (wedging) amounted to 3.6 W/kg (water temp 14°C). Minimal metabolic rate of platypuses resting on the water surface was found to be 3.9 W/kg (water temp 20°C) while minimal resting rate on land was 2.1 W/kg (air temp 26°C).

**Measurement of nectar and pollen intake
of free-ranging marsupial honey possums
(*Tarsipes rostratus*)**

S. D. Bradshaw and F. J. Bradshaw

*Department of Zoology, The University of Western Australia, Perth, WA 6907
Don.Bradshaw@uwa.edu.au*

Turnover rates of a combination of stable and radioactive isotopes (tritium, oxygen-18 and sodium-22) have been used to estimate daily intake of pollen and nectar in free-ranging honey possums in Scott National Park in the extreme southwest of Western Australia. The Field Metabolic Rate (FMR) is measured using doubly-labelled water and nectar intake is estimated independently from the turnover of both water and sodium. Nectar energy intake is then estimated from the sugar concentration of fresh nectar collected from the inflorescences on which the honey possums have been feeding. Pollen intake is finally estimated as the difference between the energy derived from nectar and the FMR. The method assumes that honey possums normally do not drink free-water in the field, that all the sodium in their diet is derived from nectar, and that individuals are in energetic balance over the measurement period. These assumptions have been tested and found to be reasonably robust, except during periods of heavy rain when some free-water intake may occur. Data from some 30 individuals show that nectar intake is reasonably constant at 6-7 ml.day⁻¹ for individuals with an average mass of 9 g. Daily pollen intake varies considerably between individuals, however, averaging 510 mg.day⁻¹ but ranging from 0 to as much as 2 g.day⁻¹. Laboratory studies with a captive colony of honey possums show that the minimum nitrogen requirement for the maintenance of balance in this species is approximately 115 mg.kg^{-0.75}.day⁻¹, well below that recorded for macropodid marsupials but very similar to that of the sugar glider, *Petaurus breviceps*, which also feeds on a high carbohydrate diet.

**Fishing for mechanisms:
the natriuretic peptide system
and the gills of the hagfish**

Will Callahan and Tes Toop

*School of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217
ttop@deakin.edu.au*

Previous studies have identified a natriuretic peptide (NP) system in the hagfish. This earlier research focussed on the Atlantic hagfish, *Myxine glutinosa*. Recently, whole gill perfusion studies and myography of afferent and efferent branchial arteries have implicated NPs in the control of blood flow through the gills of the New Zealand hagfish, *Eptatretus cirrhatus* (Forster, Glover and Simpson, XIII Int. Congress Comp. Endocrinol. 1997). However, the NP receptors that would mediate this regulation have not been identified in this species. The current study demonstrates the presence of NP binding sites in the gills of *E. cirrhatus* and examines whether NP actions in the gills of this hagfish could be mediated by guanylate cyclase receptors, similar to those found in jawed vertebrates.

**Organ system correlates with variation
in aerobic performance in red junglefowl**

M. A. Chappell and K. Hammond

*Department of Biology, University of California, Riverside, CA (USA)
chappell@citrus.ucr.edu*

Limits to whole-animal performance may be set by peripheral effectors, central organs, or by all components in synchrony ("symmorphosis"). We examined these concepts using aerobic performance in red junglefowl as a model system. We measured individual variation in the masses of peripheral (pectoralis and leg muscles) and central organs (heart, lungs, gonads, liver, spleen, and gut), and searched for correlations with minimal and maximal aerobic metabolism (BMR and VO_{2max}). We found substantial variance within and between sexes, but few consistent relationships between performance and organ size. On average, males had lower BMR and higher VO_{2max} than females, and had smaller visceral organs (gut, liver) and larger cardiovascular organs (heart, lung, haematocrit) and muscles. BMR was not correlated with any single organ mass in either sex (there was no relationship between female reproductive status and either BMR or VO_{2max} , although gonadal and oviduct mass varied more than 12-fold). In males only, BMR was correlated to combined gut mass. Male VO_{2max} residuals were correlated to heart and muscle mass; female VO_{2max} residuals were correlated to haematocrit. These results suggest that attaining high aerobic performance requires increased investment in muscles and in organs involved in oxygen delivery, but not in digestive organs.

**The effect of total water deprivation
on the natriuretic peptide system of the hopping mouse,
*Notomys alexis***

J. A. Donald, B. Davis and K. Minerds

School of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217
jdonald@deakin.edu.au

The natriuretic peptide system counters hypervolaemia and concomitant hypertension by renal diuresis and natriuresis, and vasodilation. However, the importance of hypervolaemic regulation in mammals has been questioned, and in particular, the role of natriuretic peptides in the maintenance of normal fluid balance. We are using the hopping mouse, *Notomys alexis*, to investigate the function of natriuretic peptides in a desert-adapted mammal that can survive without free access to water. Atrial natriuretic peptide (ANP) cDNA was cloned from *N. alexis*, and the sequence of ANP (1-28) was identical to rat ANP. This information enabled a radioimmunoassay for rat ANP to be used to quantify ANP levels in the heart and plasma of *N. alexis*. *N. alexis*, of either sex, were divided into an experimental group (n=12) subject to seventeen days of total water deprivation (TWD), and a control group (n=12) that had free access to water. No significant difference in haematocrit, skeletal muscle and liver water content, and cardiac ANP levels was observed between control and experimental groups. However, plasma ANP and the density of ANP binding on renal glomeruli were significantly lower in TWD animals. This data suggests that the ANP system is down-regulated during TWD, probably to conserve salt and water.

**Steroid biosynthesis in a viviparous reptile,
Tiliqua nigrolutea (Scincidae)**

A. Edwards, S. M. Jones and N. Davies

School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001
Ashley.Edwards@utas.edu.au

Predominance of either the delta-4 or delta-5 pathway for gonadal steroid biosynthesis varies between vertebrate classes. More recent information suggests that variation between both sex and reproductive condition can also occur. We examined *in vitro* biosynthesis of progesterone (P4), androstendione (AD), dehydroepiandrosterone (DHA), testosterone (T) and 17 β -oestradiol (E2) from pregnenolone (P5). Late pre-ovulatory (Oct) and post-partum (Mar) ovaries and final-spermatogenesis (Sep) and quiescent (Feb) testes of a viviparous skink, *Tiliqua nigrolutea* were used. Steroids were identified using various thin layer chromatography (TLC) solvent systems and HPLC (attached to a radiometric detector) elution times. A strong preference for the delta-4 pathway was apparent in both sexes and at both reproductive and quiescent times of year. However, for several key steroids, the relative proportions synthesised during incubation differed between reproductive conditions. Interestingly, we have demonstrated that ovarian follicles do not produce E2 from P5 *in vitro* at either time of year. The end product of P5 metabolism is instead another, more polar steroid, which is currently unidentified.

Abundance of endocrine cells in the pancreatic islets of macropods

P. Edwards, B. E. Tuch and A. M. Beal

School of Biological Science, University of New South Wales and Dept of Endocrinology, Prince of Wales Hospital, Sydney, NSW 2052
m.beal@unsw.edu.au

For more than twenty years, it has been believed that the islets of Langerhans from the macropod pancreas were essentially similar to those from the sheep pancreas in having a preponderance of α cells (glucagon-secreting) and few β cells (insulin-secreting) in comparison with islets from monogastric animals. This was argued to be an adaptation to foregut fermentation. On finding that the islets of one tammur wallaby had high β and low α cell numbers we have reassessed the proportions of the 4 cell types in islets from red and eastern grey kangaroos, tammars and sheep using immunohistological techniques. The results confirm that sheep islets have high α cell (53.9 \pm 3.20 %) and low β cell numbers (29.1 \pm 1.61 %) with the remainder being somatostatin-secreting δ cells (6.2 \pm 1.96 %) and pancreatic polypeptide secreting ϕ cells (10.8 \pm 4.04 %). In contrast, reds, greys and tammars were similar in having nearly twice as many β cells as α cells averaging 52.4 \pm 4.36 % β -cells, 28.9 \pm 2.27 % α -cells, 9.2 \pm 1.33 % δ -cells and 10.5 \pm 2.58 % ϕ -cells. Apparently, high α cell numbers are not an essential adjunct to being a foregut.

Effects of murine leptin on energetics and torpor of the dasyurid marsupial *Sminthopsis macroura*

Fritz Geiser¹, Gerhard Körtner¹ and Ingrid Schmidt²

¹Zoology, School of Biological Sciences, University of New England, Armidale, NSW 2351; ²Max-Planck-Institut für physiologische und klinische Forschung, W.G. Kerckhoff-Institut, D-61231, Bad Nauheim, Germany
fgeiser@metz.une.edu.au

Leptin plays an important role in regulating body fat stores of placental mammals, but the contribution of changes in energy uptake and expenditure to this adjustment remains controversial. We were interested in how recombinant murine leptin would affect metabolic rate and body temperature of a marsupial mammal (*Sminthopsis macroura*, 25 g) known to display daily torpor. In a group of 8 animals, food-deprived for 1 day at 18 °C, leptin treatment (5 μ g/g body mass in Tris buffer) halved the duration of torpor bouts, and raised the average daily minimum body temperature by 4.5 °C and the minimum metabolic rate by 2.2-fold in comparison to control treatment with Tris buffer. Leptin treatment thus increased daily energy expenditure by 9% although during the activity phase metabolic rates and body temperatures were not raised. Body mass was also not affected. These findings suggest that leptin affects the adjustment of thermoregulatory energy expenditure during the rest phase. They show that the hormone has strong physiological effects despite the lack of thermogenetically active brown fat in marsupials and some structural differences between murine and marsupial leptin.

The function of the vaginal caecae of the marsupial bandicoot, *Isodon macrourus*

Robert T. Gemmell and Brian V. Harmon

Department of Anatomical Sciences, University of Queensland,
St. Lucia, QLD 4072
R.Gemmell@mailbox.uq.edu.au

In the bandicoot the lateral vaginae are straight and about 50mm long and there are large lateral anterior expansions at the distal ends, called the vaginal caecae. After coitus the caecae are dilated and filled with a clear viscid semifluid containing abundant numbers of sperm. No direct function other than a store for sperm has been suggested for the caecae. In the present study the structure and function of the vaginal caecae, the longevity of sperm in the vaginal caecum and the conception rate of the bandicoot *Isodon macrourus* will be examined. Material from the lumen of the vaginal caecae obtained using a polyeththylene tube was placed on a microscope slide and the presence of motile sperm noted. Motile sperm were detected in the vaginal caecae approximately 12 hours after mating and non motile sperm were present after 18 hours. Sperm without heads were observed 6 to 8 days after conception. The vaginal caecae of the bandicoot may act as a short-term storage area that allows viable sperm to be released over several hours. These sperm fertilise the ova as they pass through the uterine tube and allow the bandicoot a high rate of conception. The conception rate for *I. macrourus* is 100% (N=8).

A comparative study of fuel choices by cells

Michael Guppy

Department of Biochemistry, University of Western Australia, Nedlands, WA 6907
mguppy@cyllene.uwa.edu.au

Modern biochemistry and physiology increasingly rely on the use of isolated cell preparations. But despite this, there are few data on what fuels cells use in an *in vitro* physiological situation to produce the ATP required to support energy-consuming processes, and no data on the comparative aspect of fuel choices by cells. This deficiency has been recently redressed by Guppy *et al.* (1997) who took advantage of the fact that human platelets can be incubated *in vitro* in their *in vivo* medium (human plasma). The data showed that lactate production and lipid oxidation accounts for 62% of the ATP turnover with minor contributions from the oxidation of glucose, ketones and glutamine, leaving 26% of the ATP turnover as yet unaccounted for. In a similar experiment using trout platelets in an artificial plasma, lactate production and fatty acid oxidation only accounted for 17% of ATP turnover, with again trivial contributions by the oxidation of glucose and glutamine. Mammalian and fish cells therefore show some marked differences in terms of fuel choices. This is interesting in itself, but also needs to be taken into account when designing culture media, if physiologically meaningful data are to be produced.

Guppy M, Abas L, Neylon C, Whisson M E, Pethick D W Niu X W. (1997) Fuel choices by human platelets in human plasma. *Eur. J. Biochem.* 244: 161-167.

Macropodids express alpha cardiac myosin in their jaw-closing muscles

J. F.Y. Hoh, Y. Kim, L. G. Sieber and C. A. Lucas

Department of Physiology, University of Sydney, Sydney, NSW 2006
joeh@physiol.usyd.edu.au

Jaw-closing muscles of vertebrates are functionally extremely diverse, having to adapt to different types of food they handle. Carnivores, primates and chiropterans express superfast myosin, rodents express fast myosins, while eutherian herbivores express slow myosin (= cardiac V3) in their jaw muscles. We investigated whether the macropodids, being herbivores, also express V3 myosin. Pyrophosphate gel electrophoresis revealed that ventricular myosin from the tamar wallaby and the red kangaroo had three ventricular myosins, V1, V2 and V3. Masseter myosin from the macropodids was homogeneous and comigrated with V1, the high ATPase cardiac isoform. An antibody raised in rabbits against red kangaroo masseter myosin reacted immunohistochemically specifically with macropodid jaw fibres, as well as atrial and ventricular muscles, but did not react with limb muscle fibres. In pyrophosphate Western blots, this antibody reacted with V1 (myosin heavy chain composition: $\alpha\alpha$) and V2 ($\alpha\beta$), but not with V3 ($\beta\beta$). These results confirm that macropodid jaw muscle express α MyHC rather than β MyHC. This difference in jaw myosin expression between eutherian herbivores and macropodids may be related to the fact that macropodids do not ruminate, and need faster and more powerful jaw muscles to comminute food only once to the correct particle size for gastric fermentation.

Thermogenesis is marsupials: Uncoupling the evidence.


P. Hope¹, R. Baudinette², H. Turnbull¹, J. Kumaratilake³ and G. Wittert¹

Departments of Medicine¹, Zoology² and Anatomy³, University of Adelaide,
Adelaide, SA 5000
gwittert@medicine.adelaide.edu.au

We have previously reported the presence of UCP1 in the dasyurid marsupial *S. crassicaudata*. Furthermore *S. crassicaudata* increases thermogenesis in response to both catecholamines and food intake, although UCP1 may not be involved in this process. In the 180 day old macropod *Macropus eugenii* (tamar wallaby), electronmicroscopy of interscapular fat showed mostly cells typical of white adipocytes; a few multilocular cells with a higher proportion of mitochondria were also seen. By contrast in perirenal fat there was a much higher proportion of these cells. While we did not detect a 32kd band consistent with UCP1 on an immunoblot, we found a much smaller band (~18kd) in the 200, but not 180 day old wallabies. Expression was greatest in perirenal fat and lowest in subcutaneous fat. Both GDP binding and thermogenesis were greater at 200 as compared to 180 days. Interestingly a ~1.2 kb mRNA band was seen in perirenal fat but not cortex, muscle or liver on an RNA blot hybridised with an oligoprobe specific for UCP1.

Recently other members of the uncoupling protein family (UCP2,3,4) have been cloned. We have identified the expression of UCP2 mRNA in both *S. crassicaudata* and *M. eugenii*, where expression is greater in perirenal than subcutaneous fat. In the new born wallaby, UCP2 expression is very low in perirenal fat and barely detectable in muscle. By 150 days expression is markedly increased in both perirenal fat and muscle, but not subcutaneous fat. No further increase in expression was seen at 200 days.

These data suggest that uncoupling proteins play a role in marsupial thermogenesis, even in the absence of clearly defined collections of brown adipose tissue.



Gonadal steroids and alterations in photoperiod affect metabolic efficiency in female *S. crassicaudata*

P. Hope¹, W. Breed², H. Turnbull¹, J. Morley¹, M. Horowitz¹ and G. Wittert¹

Departments of Medicine¹ and Anatomy², University of Adelaide, Adelaide, SA 5000.
gwittert@medicine.adelaide.edu.au

In the marsupial *S. crassicaudata*, adiposity increases in animals housed under short day photoperiods (SD), as compared to long day photoperiods (LD). This effect is seen in both sexes, and occurs without an increase in energy intake, suggesting increased metabolic efficiency. To determine the influence of gonadal steroids on the response to photoperiod, female animals were either gonadectomised or sham operated and maintained under LD or SD for 103 days. Alterations in photoperiod had no effect on body weight in the sham controls, however SD housed gonadectomised animals gained weight over the first 60 days and thereafter body weight returned to baseline. Tail width (a reflection of body fat stores) increased in both sham ($P < 0.05$) and gonadectomised ($P < 0.01$) animals under SD. However on LD, tail width increased only in the gonadectomised animals and this occurred in the first 20 days ($P < 0.01$) with no further increase thereafter. By contrast in the SD gonadectomised animals tail width increased only after day 20 ($P < 0.01$). Gonadectomy resulted in a decrease in UCP2 expression in muscle, but not fat. There was no effect of photoperiod on UCP2 expression in any tissue. These data suggest that (i) body fat stores in *S. crassicaudata* are affected by gonadectomy and alterations in photoperiod (ii) both gonadal steroid dependent and independent mechanisms mediate the effect of photoperiod (iii) alterations in UCP2 expression may account for the effect of gonadectomy, but not photoperiod and (iv) expression of UCP2 is differentially regulated in muscle and fat.

Effects of oral administration of melatonin on the reproductive cycle of a small marsupial, *Antechinus stuartii*

B. M. McAllan¹ and J. M. P. Joss²

¹ Human Biology, School of Biological Sciences, University of New England, Armidale, NSW 2351

² School of Biological Sciences, Macquarie University, North Ryde, NSW 2109
bmcallan@metz.une.edu.au & jjoss@rna.bio.mq.edu.au

Antechinus stuartii is a small marsupial with a brief, highly synchronised mating period believed to be controlled by the rate of change of photoperiod. Photoperiodic response in mammals is co-ordinated by the pineal gland and its hormone melatonin. The pineal hormone, melatonin, administered in the drinking water from the winter solstice, was found to affect the normal response of *A. stuartii* to increasing rate of change of photoperiod. Melatonin shifted the induction of estrus in the females from the first week of August (controls) to an earlier time of mid-July and the consequent pouch changes associated with pregnancy and pseudopregnancy were also shifted by the same length of time. The post-mating decline and consequent death in the males were also accelerated. The results were significantly different between control and melatonin treated groups ($p < 0.05$). The hormone melatonin appears to be important in the timing of reproduction in this marsupial.

**Osmoregulation and ionoregulation
in the Port Jackson shark,
*Heterodontus portusjacksoni***

Dennyse Newbound, Jamie O'Shea and Philip Withers

Zoology Department, The University of Western Australia, Nedlands WA 6907
dnewb@cyllene.uwa.edu.au

A comprehensive anatomical and physiological study of the Port Jackson shark, *Heterodontus portusjacksoni*, which migrates into estuarine environments to lay eggs showed relationships between the structure of the three organs involved in osmoregulation in elasmobranchs, the gills, kidneys and rectal salt gland, and how their function at lowered salinity may be achieved. Experimental acclimation to 67% seawater was undertaken over a period of six days.

The plasma of the Port Jackson shark is slightly-hyperosmotic to the environment at 100% and 67% seawater with high elevated levels of urea. The rectal salt gland has a complex structure, with the flow of secretory tubules opposite to the capillaries, in a counter-current arrangement. The counter-current organisation and the lobulate arrangement of the gland function to concentrate rectal gland fluid which is predominantly sodium, chloride and magnesium. The structure of the glomeruli in the kidney suggests an important ultrafiltration role for the nephron. The significance of this process may be replaced by secretion in lowered salinity. Chloride cells in the gills are likely to function in the elimination of ions.

The osmotic strategy used by the Port Jackson shark when adjusting to dilute media was to maintain urea concentrations at the expense of other electrolytes (especially NaCl). This strategy is suggested to be economical, as electrolytes can rapidly be elevated by the passive diffusion influx of ions when sharks return to a marine environment. It is in contrast to the general pattern in elasmobranchs that acclimation to dilute external environment is achieved by a reduction in urea concentration, with minor reductions in electrolyte concentrations.

Using dataloggers to investigate hibernation in echidnas.

S. C. Nicol and N. A. Andersen

Division of Anatomy and Physiology, University of Tasmania, GPO Box 252-24,
Hobart, TAS 7001
s.c.nicol@utas.edu

Grigg, Beard & Augee used implanted temperature sensitive radio transmitters in free ranging animals to demonstrate that echidnas are true hibernators, but temperature transmitter based systems are prone to problems, such as loss of data when animals move out of range. We have used two different types of implanted datalogger to study echidnas throughout the year: a one channel logger that recorded body temperature, and a two channel logger that recorded body temperature and activity. The activity logger had the capacity to provide detailed information on activity patterns, and while it produced some useful data it proved unreliable. The temperature loggers produced more insights into echidna physiology than had been expected: as well as demonstrating an unexpectedly early entrance into hibernation by male echidnas the temperature records could be used to clearly identify reproductive activity of females.

Grigg G C, Beard L A & Augee M L (1989). Hibernation in a monotreme, the echidna (*Tachyglossus aculeatus*). *Comp. Biochem. Physiol.* 92A: 609-612.

The innervation of the heart of the fat-tailed dunnart *Sminthopsis crassicaudata*

J. E. O'Shea

Department of Zoology, University of Western Australia, Nedlands, WA 6907
jeoshea@cyllene.uwa.edu.au

The innervation pattern of the heart of the dasyurid marsupial, *Sminthopsis crassicaudata*, by autonomic nerves was studied using isolated cardiac preparations. The atrial cardiac chambers and strips of the ventricular muscle were maintained under controlled conditions and allowed to beat spontaneously or were electrically paced. The presence or absence of autonomic nerves was demonstrated by recording any changes in the contractile force of the cardiac preparations in response to electrical stimulation across their walls. Nerve-mediated responses to the stimulation of the intramural nerves were mimicked with applied neurotransmitters and challenged by applying known neuroreceptor blocking reagents. Stimulation of intramural nerves in the ventricles caused a substantial inhibition of contractile force mediated by cholinergic nerves and an excitation mediated by adrenergic nerves. The inhibition to ventricular contractile force which occurred in response to nerve stimulation and applied acetylcholine occurred in the absence of any adrenergically maintained tone. Although the innervation pattern of the dunnart atria is typically mammalian that of the ventricles is not. The presence of a functional adrenergic innervation in the ventricles of a mammal that is able to enter and arouse from torpor is contrary to accounts for most hibernating mammals. This adrenergic ventricular innervation may be involved in the large increases in cardiac output associated with the massive adrenergic activation that mediates arousal from torpor. The occurrence of a cholinergic vagal innervation capable of inhibiting the basal force of ventricular contraction has been described in only one other group of mammals, the bats.

The down-regulation of protein synthesis during metabolic depression in the land snail *Helix aspersa*

J. L. Pakay¹, A. A. Hobbs¹, P. C. Withers² and M. Guppy¹

Departments of Biochemistry¹ and Zoology², University of
Western Australia, Nedlands, WA 6907
jlpakay@cyllene.uwa.edu.au

Metabolic depression represents a reduction in the basal metabolic rate of an animal in the order of 60-100% (1) and is employed as an adaptive mechanism in response to environmental stress. Metabolic depression is far from a rare phenomenon and has been reported in most animal phyla (2). The land snail *Helix aspersa* provides a convenient model for the study of metabolic depression as it readily enters a period of dormancy (aestivation) when deprived of food and water (in the absence of any temperature change). Associated with this dormancy is a high degree of metabolic depression, an 84% decrease in metabolic rate as measured by oxygen consumption (3). Protein synthesis is a major contributor to cell energy expenditure and thus is thought to be one of the major energy utilising processes which must be down-regulated during metabolic depression. The aim of this study was to measure the rate of protein synthesis in active and aestivating snails in two major tissues *in vivo*, foot muscle and hepatopancreas (digestive gland). It was found that the rate of protein synthesis decreased by 86% and 82% respectively. Evidence from other systems suggests that the down-regulation of protein synthesis during metabolic depression is controlled at the level of initiation of translation. One of the most well characterised mechanisms of translational control is down-regulation by phosphorylation of the alpha subunit of eIF-2 (eukaryotic initiation factor 2). Therefore the phosphorylation state of eIF-2alpha is being investigated as well as kinase activity towards eIF-2alpha during both the active and aestivating states.

(1) Guppy *et al.* (1994). *Comp. Biochem. Physiol.* 109 B (2/3): 175-189; Guppy M. & Withers P. (1999) *Metabolic depression in animals: Physiological perspectives and biochemical generalisations*. *Biol. Rev. in press*; (3) Pedler *et al.* (1996). *J. Comp. Physiol. B* 166: 375-381.

Physiological precocity in an altricial parrot

J. T. Pearson

Department of Electrical and Electronic Engineering, Muroran Institute of Technology, Muroran, Mizumoto-cho 27-1 JAPAN
jpearson@muroran-it.ac.jp

Parrots lay small eggs and hatch with the lowest degree of morphological and behavioural maturity. Previously one parrot species, *Agapornis roseicollis*, was shown to have a low degree of functional maturity during both the embryonic and posthatching periods. In this study the cockatiel, *Nymphicus hollandicus*, was found to have low functional maturity during incubation, i.e. a low hatchling dry mass fraction and low metabolic rates of embryos and hatchlings relative to that of predicted conspecific adults of similar masses, which is similar to most altricial birds. However, in spite of an absence of natal down, cockatiel chicks develop strong thermogenic powers within 4-5 d at small body masses, exceeding that of predicted adults at the same mass, much earlier in development than in other altricial species. The development of posthatching metabolism in this parrot resembles the patterns of small precocial species, emphasising that there is more diversity among altricial birds than recognised in even the most recent studies, and that there is continuous variation in developmental modes with regard to physiological maturity.

Ingestion of extract of leaves of mangosteen (*Garcinia mangostana* Linn) induces germ cell apoptosis in rat testis

Y. V. A. Riono¹, D. Blache¹, A. M. Dharmarajan² and G. B. Martin¹

Faculty of Agriculture (Animal Science)¹ and Department of Anatomy and Human Biology², University of Western Australia, Nedlands, WA 6907
ariono@cyllene.uwa.edu.au

Indonesian women drink tea made from mangosteen as a contraceptive. This herbal medicine might also work in males because it decreases sperm production in rats (Riono *et al.*, unpublished). However, the mechanism is not clear. We tested whether mangosteen extract reduces germ cell numbers by promoting apoptosis, the controlled cell death that is characterized by cellular shrinkage, chromatin condensation, membrane blebbing and orderly DNA fragmentation. This process is thought to explain 75% of the normal loss of mature sperm cells in adult mammalian testis. Male rats were given either a control (vehicle) or treatment (a daily oral dose of a double methanol extract of mangosteen leaves, 100 mg/kg BWt) for 8 weeks. Five rats from each group were killed every 2 weeks and testicular tissue was removed and subjected to *in situ* 3'-end-labeling of internucleosomal DNA fragments. Apoptosis increased in extract-treated rats after 4 weeks of treatment, suggesting that reduction of sperm production by mangosteen extract is due to an increase in apoptosis, probably in the germ cells themselves. Further studies are needed to elucidate the active components of the extract and to test whether the extract directly induces apoptosis, or whether other factors are involved.

Ecological correlates of locomotor performance in geckos

Alan R. Roberts

Department of Zoology, University of Western Australia, Nedlands, WA 6907
aroberts@cyllene.uwa.edu.au

In many diurnal lizards, aspects of locomotor performance such as endurance and sprint speed, and their thermal dependencies, are often closely related to ecological or physiological traits such as foraging mode and the preferred body temperature for activity. In the few nocturnal lizards examined to-date, however, sprint performance seems to peak at body temperatures well above those experienced during routine activity and endurance capacities that are often assumed to be limiting. To further investigate these apparent differences, an examination of the inter-specific differences in endurance and sprint speed was undertaken for 5 species of sympatric nocturnal gecko and a diurnal agamid from the semi-arid region of Western Australia. Maximum sprint speeds and endurance times varied dramatically between the five species of gecko. Species thought to be exposed to the greatest risk of predation had the fastest sprint speeds, whereas those that moved most frequently while foraging had the greatest endurance. Body temperatures optimal for sprinting were higher in all geckos than those optimal for endurance. Peak sprint performance appeared to occur at body temperatures typically encountered during times of the greatest risk of predation i.e. during the daytime. Endurance peaked at the lower body temperatures typically experienced during nocturnal activity. In contrast to the geckos, sprint speed and endurance had similar thermal optima in the diurnal agamid, both coinciding with its daytime activity temperatures. In all species, sprint speed had a wider thermal performance breadth than endurance, possibly reflecting the importance of being able to escape from potential predators regardless of body temperature.

Aldosterone and renal function in avian species

Julie Roberts

Division of Animal Physiology, School of Rural Science and Natural Resources, University of New England, Armidale, NSW 2351
jroberts@metz.une.edu.au

The role of aldosterone in avian kidney function has not been clearly demonstrated although the assumption has been that the action of aldosterone in avian species is similar to that in mammals. Ten adult roosters were used to investigate the effect of aldosterone infusion on plasma electrolyte composition and renal function. Birds were anaesthetised with DIAL and a carotid artery and superficial ulnar vein cannulated. The hindgut was blocked off and a tube inserted for collection of ureteral urine. The five birds comprising the control group (CONT) were infused at 0.2 mL/min/kg with a solution of isotonic saline containing 10 g/kg inulin and 10 g/kg PAH, for a total of 8.5 hours. The five roosters in the aldosterone treatment group (ALDO) were infused with the same solution except that aldosterone (125 ug/L) was added to the infusate after the initial 30 minutes of the experiment. Blood samples were collected hourly and urine samples every 15 minutes. Plasma and urine samples were analysed for inulin, PAH, osmolality, sodium (Na), potassium (K), chloride (Cl). Plasma samples were taken for aldosterone assay at four stages of the experiment. The plasma aldosterone concentrations achieved represented a high physiological level. Plasma K increased significantly, urine K was significantly lower and the urine:plasma ratio for K was lower in the ALDO group. An increased urine flow rate in the ALDO group was accompanied by lower urine osmolality and chloride and lower urine:plasma ratio for osmolality. Osmolar clearance and total osmolar output were significantly higher for the ALDO group. The total excretion rate and the fractional excretion of sodium were higher in the ALDO group throughout the experiment and there was a transient decrease about 45 minutes after the start of the aldosterone infusion. Glomerular filtration rate (GFR) was not significantly affected by aldosterone infusion whereas the clearance of PAH (C_{pah} - an indication of renal plasma flow) remained constant in the ALDO group at the same time as it increased in the CONT group. This resulted in a higher ratio of GFR:C_{pah} in the ALDO group. These results indicate that aldosterone has direct effects on the avian kidney but these effects differ from those in mammals.

**Effect of aerial exposure on the western rock lobster,
*Panulirus cygnus***

Patrick T. Spanoghe, Glen W. Davidson and Brian D. Paterson

School of Biomedical Sciences, Curtin University of Technology,
GPO Box 1987 Perth, WA 6001
rspanogh@cc.curtin.edu.au

Post-harvest operations play a critical role in determining the quality and value of live seafood products, such as the western rock lobster, *Panulirus cygnus*. Although lobsters should be kept in water as much as possible after harvest, lobsters do experience unavoidable episodes of emersion, as may occur during transport to processing facilities or to domestic or foreign markets. Given these considerations, this study sought to examine the effect of a 24 h period of emersion followed by a 48 h period of reimmersion, at 20°C, on the physiology of western rock lobsters.

The results revealed that, when exposed to air, both pre- and post-branchial PO₂ decreased and pre-branchial PCO₂ increased, indicating that gill function was compromised. A progressive increase in haemolymph lactate titre during emersion suggested an increased dependence on anaerobic metabolism when in air. The increase in lactate concentration and PCO₂ led to a mixed respiratory/metabolic acidosis and a fall in haemolymph pH. A classic hyperglycaemia was also observed. The results suggest that although a depression of behavioural responses to handling could be detected during emersion and following re-immersion, western rock lobsters can defend tissue energy levels anaerobically, with no change in adenylate energy charge (AEC) of abdominal muscle being recorded. Perturbations in the levels of potassium, magnesium and calcium in the haemolymph were recorded during both emersion and following re-immersion. Recovery of all parameters was complete within 8 h after re-immersion.

**A cholinergically mediated adrenergic response
in the heart of the horned shark,
*Heterodontus portusjacksoni***

Anton Thompson and Jamie E. O'Shea

Department of Zoology, University of Western Australia, Nedlands, WA 6907
apt@cyllene.uwa.edu.au

The atypical excitatory effect of acetylcholine (Ach) on cardiac function was investigated in the horned shark, *Heterodontus portusjacksoni*. Electrically paced ventricular strips showed a striking 391 ± 26% increase on basal force of contraction as a response to exogenously applied Ach. The response was similar in nature to the response of electrically paced ventricular strips to adrenaline (Ad) which caused a 383 ± 72% increase. The response to Ad was faster than the response to Ach, especially over the first 10 mins after application. The response to Ach was blocked by the muscarinic antagonist, atropine and the β-adrenergic antagonist, propranolol, and was reduced by bretylium, an agent known to inhibit the release of catecholamine from adrenergic nerves. The evidence from this study suggests that Ach caused a localised release of a catecholamine via muscarinic receptors. A cholinergic-adreno complex has been proposed suggesting that elasmobranchs may be capable of finer control of cardiac output than was previously suspected. Investigations on the spontaneously beating atrium gave results in accordance with previous authors with no evidence of an excitatory response. Exogenously applied Ach caused the atrium to beat slower and less forcibly. Atropine blocked the response before and after it was initiated suggesting Ach was effecting the muscle directly via muscarinic receptors.

Comparing the metabolism of reptiles with varying body mass

G. G. Thompson

Centre for Ecosystem Management, Edith Cowan University,
100 Joondalup Drive, Joondalup, WA 6027
G.Thompson@cowan.edu.au

How useful are the generally accepted allometric exponents of 0.75 (interspecific) and 0.67 (intraspecific) when comparing the metabolism of reptiles of varying body mass? This paper will explore some of the problems and issues of comparing the standard metabolic rate of reptiles (specifically goannas and pythons) of different body mass. It will look at species which have an intraspecific mass exponent not approximately 0.67, genera where the interspecific mass exponent is not 0.75 and species where neonates don't appear to conform to the intraspecific allometric relationship evident for adults. It will also address some related statistical issues.

The MO_2 and metabolic release of CO_2 from labeled glucose and lactate in the exercising ghost crab, *Ocypode ceratophthalmus*

W. J. van Aardt

Zoology Department, Potchefstroom Universiteit, Private Bag X6001,
Potchefstroom, South Africa 2520
drklva@puknet.puk.ac.za

It is known that in ghost crabs the gills act as osmoregulatory organs and do not participate in the CO_2 and O_2 exchange process during respiration. To determine the efficiency of the relatively thin branchial lining as a functioning "lung", crabs were run on a treadmill for 20 minutes. During exercise the strong and weak points of the respiratory physiology of the crab could be determined. Before the run each crab was injected with radioactively labeled glucose or l-lactate. It was found that haemolymph l-lactic acid levels increased to 20 mmol L⁻¹; the heart rate increased from 100 beats per minute to nearly 400 beats per minute; MO_2 increased about five times above resting levels. The oxygen dissociation curves constructed for the blood from resting crabs and exercising individuals shows a large shift to the right. A linear increase of $^{14}CO_2$ from injected l-lactate indicates that this gas is nearly instantaneously released by the branchial lining or "lung". The same result was found when CO_2 was metabolically released from injected glucose. These findings, together with the MO_2 measurements gave evidence that the branchial lining has a gas exchange capacity similar to that of an ectothermic vertebrate.

Gonadal hormones are proximate factors controlling seasonal fat metabolism in adult male emus.

J. Van Cleeff, D. Blache, I. Malecki and G. B. Martin

Animal Science Department, University of Western Australia, Nedlands, WA 6907
jvanclee@agric.uwa.edu.au

Mature male emus fed *ad libitum* show inversely related seasonal patterns of liveweight and sex hormones. Feed intake is lowest in winter but doubles by 5 weeks after the winter solstice, and reaches a maximum near the spring equinox. Liveweight is highest in summer, lowest 5 weeks after the winter solstice, but returns to summer values within 3 weeks of the spring equinox. Testosterone is detectable from 9 weeks before until 7 weeks after the winter solstice.

Mature male emus that had been castrated at 8 weeks of age ate similar amounts of feed to intact males in spring, summer and autumn. In winter, intake was maximum one week after the solstice and higher than for intact birds. Liveweight of castrated emus followed a similar pattern to intact birds, except that minimum values were observed in the week of the winter solstice. Castrates weighed less than intact birds, except in winter.

Early castration attenuated vernal fattening of male emus, while appetite was only slightly affected. The pattern of feed intake and fat gain and loss was shifted to earlier in the year for castrates. These results confirm for emus the interaction between photoperiod and gonadal steroids in the control of fat deposition and metabolism.

Prostaglandin $F_{2\alpha}$ receptors in the brushtail possum, *Trichosurus vulpecula* and the short-nosed bandicoot, *Isodon macrourus*, at birth.

Colleen E. Veitch¹, Conrad Sernia² and Robert T. Gemmell¹

Departments of Anatomical Sciences¹ and Physiology and Pharmacology²,
University of Queensland, St Lucia, QLD 4072
C.Veitch@mailbox.uq.edu.au

Prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) is involved with the regression of the corpus luteum (CL). Regression occurs at parturition in the possum but later in the bandicoot. The plasma concentration of 13, 14-dihydro-15-keto-prostaglandin $F_{2\alpha}$ (PGFM), the major metabolite of $PGF_{2\alpha}$, is briefly elevated at birth in the possum and elevated for several days around birth in the bandicoot. In both species, $PGF_{2\alpha}$ injection induces adoption of the birth position. This study investigates $PGF_{2\alpha}$ receptors in the uterus and CL of possums and bandicoots at parturition. $PGF_{2\alpha}$ receptors are not present in the bandicoot and possum uteri and the bandicoot CL. $PGF_{2\alpha}$ does not appear to be involved with uterine contractions in the possum or the bandicoot at parturition. However, low concentrations of receptors were found in the possum CL, suggesting the involvement of $PGF_{2\alpha}$ in CL regression. In the bandicoot, there were no $PGF_{2\alpha}$ receptors in the CL at birth. This provides an explanation for the failure of $PGF_{2\alpha}$ administration to regress the bandicoot CL at birth. However, $PGF_{2\alpha}$ injections will cause regression of the CL at day 30 of lactation and therefore future studies will be directed to this post-natal period.

Overview of metabolic depression in ectotherms and endotherms

Philip C. Withers

Department of Zoology, University of Western Australia, Nedlands WA 6907
pwithers@cyllene.uwa.edu.au

Metabolic depression has been recorded for virtually all major animal phyla in response to environmental stress. An allometric analysis suggests three general patterns of metabolic depression. Firstly, a common pattern is metabolic depression to approximately 0.05-0.4 of rest e.g. molluscs, earthworms, crustaceans, fishes, amphibians, reptiles. This "intrinsic" depression is in anticipation of adverse environmental conditions but without substantial changes to their state of body water. Some animals survive anoxia for limited periods, with a similar depression to 0.05-0.4. The profound decrease in metabolic rate of torpid mammals and birds is mainly due to readjustment of thermoregulatory control and a decrease in body temperature with a concomitant Q_{10} effect, although there may be a modest intrinsic metabolic depression for deep torpor (to approximately 0.6). Secondly, a more extreme pattern of metabolic depression to <0.05 of rest is evident for cryptobiotic (anhydrobiotic, osmobiotic, or cryptobiotic) animals with a profound change in their ionic/osmotic balance or state of body water. A few normally-aerobic animals can have an anaerobic metabolic rate than 0.005 of resting. The metabolic rate of anhydrobiotic animals can be unmeasurable, if not zero. Thirdly, a pattern of indefinite absence of metabolism when anoxybiotic but normally hydrated is apparently unique to diapaused eggs of the brine-shrimp (*Artemia* spp., an anostracan crustacean).

Metabolic physiology of the marsupial mole (*Notoryctes caurinus*)

Philip C. Withers¹, Graham G. Thompson² and Roger S. Seymour³

¹Department of Zoology, University of Western Australia, Nedlands, WA 6907;


²Centre for Ecosystem Management, Edith Cowan University, Joondalup, WA 6027;

³Department of Zoology, University of Adelaide, Adelaide, SA 5005

pwithers@cyllene.uwa.edu.au

The marsupial mole (*Notoryctes*) is the most unusual of all Australian marsupials, essentially living entirely underground. It closely resembles the golden moles (placental family Chrysochloridae) such as the Namib Desert golden mole (*Eremitalpa grantii namibensis*), being an excellent example of convergence in morphology. Not only is the marsupial mole unusual, it is extremely rare and nothing is known of its physiology, not even its body temperature. The essentially completely fossorial habit of the marsupial mole, in contrast with the nocturnal surface-feeding of the Namib Desert golden mole, makes the marsupial mole extremely difficult to locate for study.

We report that the marsupial mole (*N. caurinus*) has a low and labile body temperature, as expected for a fossorial mammal, but does not have a substantially lower-than-predicted BMR. Thermal conductance is high, and variable. The metabolic cost of burrowing through dry sand is very high, as expected, compared to the metabolic cost of terrestrial (running) locomotion. The kidney of the marsupial mole has only a moderate medullary area, suggesting a mesic- rather than arid-adapted water balance, reflecting the high water content of the food. With respect to its physiology, the marsupial mole is also remarkably convergent with the Namib Desert golden mole.



POSTERS

Corticosterone response of captive northern brown kiwi (*Apteryx australis mantelli*) to an acute stressorD. C. Adams¹, J. F. Cockrem¹ and M. A. Potter²¹Conservation Endocrinology Research Group, Institute of Veterinary, Animal and Biomedical Sciences; ²Institute of Natural Resources (Ecology), Massey University, Palmerston North, New Zealand.

D.C.Adams@massey.ac.nz

Corticosterone is the main adrenocortical hormone released during stress in birds, and plasma corticosterone levels usually rise rapidly after a bird experiences a stressor. In this study we examined the corticosterone response of six captive northern brown kiwi to an acute stressor (handling) by collecting blood samples from kiwi 3 times over one hour and then once more, either 3, 5 or 7 hours after the kiwi had initially been handled. All of the kiwi responded to the acute stressor with a rise in corticosterone levels. Plasma corticosterone levels increased markedly to a peak 60 min after handling then declined and remained above initial levels from 3 to 7 hours after handling. Corticosterone levels were higher than usual for birds. Results from this preliminary study also suggest that there may be a difference between the corticosterone responses of nocturnal housed and outdoor housed captive kiwi.

The use of dataloggers to determine behavioural activity in the platypus (*Ornithorhynchus anatinus*)

Philip Bethge, Helen Otley, Sarah A. Munks and Stewart Nicol

Division of Anatomy and Physiology, University of Tasmania, GPO Box 252-24, Hobart, TAS 7001
p_bethge@utas.edu.au

Studies into the behavioural activity pattern of the platypus have been restricted in the past on visual observation or radio tracking techniques. To examine platypus behaviour in the wild more closely, in this study animals were equipped with dataloggers (Minimitter Co., Inc., USA) measuring activity in epoch lengths from 2 sec to 5 min by using an accelerometer sensitive to <.05 g, integrating degree and intensity of motion. Over the last 9 months 18 individuals have been equipped with combined datalogger-transmitter packages (weight 40 g) in a subalpine lake in northwest Tasmania. Animals were recaptured and data retrieved after three to five weeks. Preliminary results indicate a high variability in activity. About 40% of animals foraged preferentially over the day while 60% preferred night time. 50% of all platypuses showed irregular activity patterns with some animals changing start and end time of their active period within days and foraging up to 20 hours without resting. On average foraging trips lasted for 13 hours daily. Mean dive times of three individuals were between 12 and 66 seconds with 47% of all dives lasting between 30 to 40 seconds. Animals performed up to 800 dives per day.

Intracerebral infusion of recombinant leptin inhibits LH secretion in male sheep by decreasing feed intakeP. Celi¹, D. Blache¹, G. B. Martin¹, P. E. Vercoc¹, R. A. Dynes² and R. Tellam³¹Faculty of Agriculture (Animal Science), The University of Western Australia, Nedlands, WA 6907; ²CSIRO Animal Production, Wembley, WA 6014; ³CSIRO

Tropical Agriculture, Indooroopilly, QLD 4068

pceli@agric.uwa.edu.au

We have previously observed that infusion of leptin reduces both LH pulse frequency and feed intake in mature Merino rams. We therefore infused bovine leptin into the third cerebral ventricle of rams fed *ad libitum* to investigate whether the inhibition of LH secretion is a direct effect of leptin or is due to a decrease in the availability of energy substrates. Seventeen rams were randomly allocated to treatment with artificial cerebrospinal fluid (aCSF) or 0.4 µg/h bovine leptin in

aCSF. The effect of decreased food intake on LH secretion was controlled by pair-feeding a third group to the leptin-infused group. The infusions were continuous for 24 h for five days. Blood plasma sampled every 20 min for 24 h on days 1 and 5 of the treatment period was assayed for LH. Leptin infusion decreased feed intake compared to the control group ($P < 0.05$). Five days of icv leptin infusion and pair-feeding decreased LH pulse frequency compared to control values ($P < 0.01$). Leptin decrease feed intake and thus energy substrates, and this reduces LH secretion. It seems unlikely that leptin directly inhibits reproduction.

Myofibrillar protein composition of regenerating and pristine claw closer muscle of the yabby, *Cherax albidus*.

L. Cutler, A. E. Patak and J. M. West

School of Natural Sciences, Edith Cowan University, 2 Bradford Street, WA 6050
a.patak@cowan.edu.au

Yabbies have the ability to regenerate lost limbs. This involves many developing stages until the new limb becomes morphologically and physiologically virtually indistinguishable from its predecessor. In the yabby, the claw becomes functional after the first moult when the dactyl becomes moveable. However, the muscle fibres appear to be at least partly functional already before this (West *et al.*, 1995). Crustacean muscles are composed of two major fibre types that differ in ultrastructure, contractile properties and protein composition. Fast fibres have short sarcomeres and low fatigue resistance, whereas slow fibres have long sarcomeres and high fatigue resistance. In addition, each fibre type has a unique myofibrillar protein profile. Muscle fibres from regenerating claws of *Cherax destructor* have previously been characterised by ultrastructure, sarcomere length and activation properties (West *et al.*, 1995). Muscle fibres isolated from regenerating claws with a fused dactyl had ultrastructural and activation properties characteristic of fast fibres with a sarcomere length typical of slow fibres. In the present study, the myofibrillar protein compositions of pristine and regenerating claw muscle fibres of the yabby were used to document muscle fibre regeneration and to compare pristine with regenerated muscle fibres.

West J. M., Humphries D. C., & Stephenson G. (1995) Characterisation of ultrastructural and contractile activation properties of crustacean (*Cherax destructor*) muscle fibres during claw regeneration and moulting. *J. Muscle Res. Cell. Motil.* 16, 267-284.


Anaerobic metabolism in the intertidal elephant snail, *Scutus antipodes* (Fissurellidae)

J. P. Elias and J. Baldwin

Department of Biological Sciences, Monash University, Clayton, VIC 3168
John.Elias@sci.monash.edu.au

The elephant snail, *Scutus antipodes*, is a common gastropod of the intertidal zone on south-eastern Australian rocky shores. It occupies a similar ecological niche to the abalone (Haliotidae) with which it is grouped in the Archaeogastropoda. Abalone display well developed capacities for anaerobic metabolism during both exercise induced and environmental hypoxia, producing D-Lactate and the unusual pyruvate reductase end product tauropine. Given the phylogenetic, ecological and behavioural similarities of *S. antipodes* and abalone it was considered of interest to compare their anaerobic capabilities.

Tauropine dehydrogenase was the only pyruvate reductase enzyme detected in *S. antipodes* foot muscle, and tauropine accumulated during both air exposure (2-6 h) and enforced exercise (shell righting behaviour). These hypoxic stresses were accompanied by decreased arginine phosphate reserves. In comparison with abalone, the lower pyruvate reductase activities and intracellular pH buffering of *S. antipodes* foot musculature imply a more limited capacity for anaerobic metabolism in these hypoxic situations.



An investigation of the factors causing the population decline of the swift parrot

B. D. Gartrell¹, S. M. Jones¹, R. N. Brereton² and L. B. Astheimer³

¹School of Zoology, University of Tasmania, GPO Box 252-05, Hobart TAS 7001; ²Department of Primary Industries, Water and Environment, Hobart TAS 7001; ³Department of Biomedical Science, University of Wollongong, Wollongong, NSW 2522
S.M.Jones@utas.edu.au

The swift parrot (*Lathamus discolor*) is a small migratory parrot, which breeds in the eucalypt forests of Southeast Tasmania. Its status has been recommended to be upgraded to "endangered" as there are less than 1000 breeding pairs in the wild. The main reasons for its decline include fragmentation and loss of foraging and nesting habitat, fluctuations in breeding success associated with years of poor flowering of the blue gum (*Eucalyptus globulus*) and finally, increased collision mortality.

This poster outlines a doctoral project with the aim to investigate those factors, which are suspected to be responsible for the swift parrot population decline. The importance of understanding the mechanisms of population decline in regard to conservation efforts and the rationale of the research plan as it relates to these mechanisms are outlined. At the conclusion of the project, the goal is to have the first detailed study of the reproductive biology and endocrinology of a threatened parrot in Australia. This will support the goals of the Swift Parrot Recovery Plan, to increase the breeding population and to preserve appropriate habitat of the swift parrot. In a broader context the project will develop investigative techniques with importance to other avian conservation programmes.

Energy expenditure and water uptake in free-ranging and captive blossom-bats, *Syconycteris australis* (Megachiroptera)

Fritz Geiser and Dionne K. Coburn

Zoology, School of Biological Sciences, University of New England,
Armidale NSW 2351
fgeiser@metz.unc.edu.au

Blossom-bats, *Syconycteris australis* (18 g) are known to be extremely active throughout the night. Since they frequently enter daily torpor, we wondered whether this may be related to high energy expenditure in the field. We measured field metabolic rates (FMR) of bats at a subtropical site using the doubly labelled water method. Captive bats at constant ambient temperature (T_a) and with food ad libitum were measured for comparison. The FMR of *S. australis* of 8.55 ml $\text{CO}_2/\text{g}\cdot\text{h}$ or 76.87 kJ/d (7.04-fold of basal metabolic rate, BMR) was one of the highest values measured in endotherms to date and may be why these bats frequently enter torpor when at rest. Daily water uptake in the field was about 1.8-fold of body mass. Use of energy in captive bats was about $\frac{2}{3}$ that of FMR. This suggests that foraging times in the field and laboratory were similar, and that daily energy expenditure was not strongly affected by T_a or food availability. Our study shows that *S. australis* has a FMR that is about 2-fold of that predicted for its size although their BMR is lower than predicted. Thus, extrapolations from BMR about energy requirements in free-ranging animals must be made with caution.

Effect of temperature and pH on motility and velocity of emu spermatozoa

Lena Holm¹, Irek A. Malecki² and Graeme B. Martin¹

¹Department of Animal Physiology, Swedish University of Agricultural Sciences, Uppsala, Sweden; ²Faculty of Agriculture (Animal Science), University of Western Australia, Nedlands, WA 6907

imalecki@agric.uwa.edu.au

Chicken spermatozoa need to be motile to reach the sperm storage tubules in the oviduct and to leave the tubules to fertilise the eggs. During storage, however, the spermatozoa are immotile. The mechanisms responsible for inactivation and

reactivation are not clear but a change in pH has been suggested, since only small changes in pH are required to stimulate or inhibit the motility of spermatozoa from several avian species. The effect of pH on emu spermatozoa was studied in semen from 4 males. Velocity ($\mu\text{m/s}$) and % motile spermatozoa were measured using digitised image analysis (about 100 sperm per experimental condition) in four media: MES (pH 6.0), TES (pH 7.0), TRIZMA (pH 8.0), and CAPSO (pH 9.0). At 20°C, motility was high in all media (pH 6.0-pH9.0) but the velocity was low. At 40°C, spermatozoa showed high motility and high velocity at pH 6.0 - pH 8.0. At pH 9.0, however, both measures were markedly reduced, although the sperm remained motile. These observations contrast with those for spermatozoa from the chicken, turkey and quail, which are inhibited at 40°C and pH 6.0-7.0 but stimulated at pH 9.0. The optimal pH range for spermatozoa may differ between the emu and other species.

The effects of moulting on muscle fibre characteristics of the yabby, *Cherax albidus*

T. Lamey and A.E. Patak

School of Natural Sciences, Edith Cowan University, 2 Bradford Street,
Joondalup, WA 6050
a.patak@cowan.edu.au

Moulting has a profound effect on the growth of crustaceans. Growth is cyclic with relatively long times of physiological stasis punctuated by short periods of rapid growth at moulting. The claw muscle is particularly affected. It undergoes extensive atrophy (up to 60% of muscle mass) to enable withdrawal through the narrow base whilst remaining functional. In contrast, the large abdominal muscle does not undergo atrophy during moulting. Crustacean muscles have two major fibre types that have previously been identified in *Cherax destructor* claw muscle; slow contracting fibres with long sarcomeres, and fast contracting fibres with short sarcomeres. All muscle fibre types are involved in moult-induced atrophy as well as moulting-associated fibre elongation, although different mechanisms appear to operate in each fibre type. In the present study, histochemical staining techniques have been developed to demonstrate contractile speed, fatigue resistance and fibre size of yabby muscle fibres. These techniques were used to determine the occurrence and distribution of fibre types and sub-types in the claw and abdomen during intermoult and to compare the fibre characteristics of moult-induced atrophic claw muscle with non-atrophic abdominal muscle over the moult cycle.

Respiratory responses of intertidal crab eggs to changes in oxygen tensions

N. Leelapiyanart¹ and H. H. Taylor²

¹Department of Aquatic Science, Burapha University, Chonburi, Thailand;

²Department of Zoology, University of Canterbury, Christchurch, New Zealand
nongnud@bucc4.buu.ac.th

In response to declining oxygen tension, pre-hatching eggs of *H. rotundifrons* and *C. lavauxi* exhibited characteristics of oxygen conformers and regulators respectively. The P_{crit} value for late stage eggs of *C. lavauxi* was approximately 6 kPa (45 mm Hg). The lactate concentration in eggs of *H. rotundifrons* increased from about 0.602 mmol.l⁻¹ in normoxic eggs to about 10.20 mmol.l⁻¹ after 3h hypoxic exposure. The rate of lactate accumulation during hypoxia suggests that the eggs are not sustaining metabolism by anaerobiosis, but are allowing metabolic depression. Pre-hatching eggs of *H. rotundifrons* and *C. lavauxi* exhibit no ventilatory or locomotory movements, and limited circulatory function. We hypothesize that the greater dependence of oxygen uptake on ambient PO_2 in the larger *H. rotundifrons* eggs reflects a greater diffusion limitation. This interpretation is supported by calculations of PO_2 gradients in a simplified model. This model showed that the centre of *Heterozius* eggs becomes anoxic at quite moderate external hypoxia, thus limiting total oxygen uptake. For *Cyclograpsus*, the centre does not become anoxic until severe external hypoxia prevails.

An investigation of the factors causing the population decline of the swift parrot

B. D. Gartrell¹, S. M. Jones¹, R. N. Brereton² and L. B. Astheimer³

¹School of Zoology, University of Tasmania, GPO Box 252-05, Hobart TAS 7001; ²Department of Primary Industries, Water and Environment, Hobart TAS 7001; ³Department of Biomedical Science, University of Wollongong, Wollongong, NSW 2522
S.M.Jones@utas.edu.au

The swift parrot (*Lathamus discolor*) is a small migratory parrot, which breeds in the eucalypt forests of Southeast Tasmania. Its status has been recommended to be upgraded to "endangered" as there are less than 1000 breeding pairs in the wild. The main reasons for its decline include fragmentation and loss of foraging and nesting habitat, fluctuations in breeding success associated with years of poor flowering of the blue gum (*Eucalyptus globulus*) and finally, increased collision mortality.

This poster outlines a doctoral project with the aim to investigate those factors, which are suspected to be responsible for the swift parrot population decline. The importance of understanding the mechanisms of population decline in regard to conservation efforts and the rationale of the research plan as it relates to these mechanisms are outlined. At the conclusion of the project, the goal is to have the first detailed study of the reproductive biology and endocrinology of a threatened parrot in Australia. This will support the goals of the Swift Parrot Recovery Plan, to increase the breeding population and to preserve appropriate habitat of the swift parrot. In a broader context the project will develop investigative techniques with importance to other avian conservation programmes.

Energy expenditure and water uptake in free-ranging and captive blossom-bats, *Syconycteris australis* (Megachiroptera)

Fritz Geiser and Dionne K. Coburn

Zoology, School of Biological Sciences, University of New England,
Armidale NSW 2351
fgeiser@metz.une.edu.au

Blossom-bats, *Syconycteris australis* (18 g) are known to be extremely active throughout the night. Since they frequently enter daily torpor, we wondered whether this may be related to high energy expenditure in the field. We measured field metabolic rates (FMR) of bats at a subtropical site using the doubly labelled water method. Captive bats at constant ambient temperature (T_a) and with food ad libitum were measured for comparison. The FMR of *S. australis* of 8.55 ml CO₂/g.h or 76.87 kJ/d (7.04-fold of basal metabolic rate, BMR) was one of the highest values measured in endotherms to date and may be why these bats frequently enter torpor when at rest. Daily water uptake in the field was about 1.8-fold of body mass. Use of energy in captive bats was about 1/3 that of FMR. This suggests that foraging times in the field and laboratory were similar, and that daily energy expenditure was not strongly affected by T_a or food availability. Our study shows that *S. australis* has a FMR that is about 2-fold of that predicted for its size although their BMR is lower than predicted. Thus, extrapolations from BMR about energy requirements in free-ranging animals must be made with caution.


Effect of temperature and pH on motility and velocity of emu spermatozoa

Lena Holm¹, Irek A. Malecki² and Graeme B. Martin²

¹Department of Animal Physiology, Swedish University of Agricultural Sciences, Uppsala, Sweden; ²Faculty of Agriculture (Animal Science), University of Western Australia, Nedlands, WA 6907

imalecki@agric.uwa.edu.au

Chicken spermatozoa need to be motile to reach the sperm storage tubules in the oviduct and to leave the tubules to fertilise the eggs. During storage, however, the spermatozoa are immotile. The mechanisms responsible for inactivation and



reactivation are not clear but a change in pH has been suggested, since only small changes in pH are required to stimulate or inhibit the motility of spermatozoa from several avian species. The effect of pH on emu spermatozoa was studied in semen from 4 males. Velocity ($\mu\text{m/s}$) and % motile spermatozoa were measured using digitised image analysis (about 100 sperm per experimental condition) in four media: MES (pH 6.0), TES (pH 7.0), TRIZMA (pH 8.0), and CAPSO (pH 9.0). At 20°C, motility was high in all media (pH 6.0-pH9.0) but the velocity was low. At 40°C, spermatozoa showed high motility and high velocity at pH 6.0 - pH 8.0. At pH 9.0, however, both measures were markedly reduced, although the sperm remained motile. These observations contrast with those for spermatozoa from the chicken, turkey and quail, which are inhibited at 40°C and pH 6.0-7.0 but stimulated at pH 9.0. The optimal pH range for spermatozoa may differ between the emu and other species.

The effects of moulting on muscle fibre characteristics of the yabby, *Cherax albidus*

T. Lamey and A.E. Patak

School of Natural Sciences, Edith Cowan University, 2 Bradford Street,
Joondalup, WA 6050
a.patak@cowan.edu.au

Moulting has a profound effect on the growth of crustaceans. Growth is cyclic with relatively long times of physiological stasis punctuated by short periods of rapid growth at moulting. The claw muscle is particularly affected. It undergoes extensive atrophy (up to 60% of muscle mass) to enable withdrawal through the narrow base whilst remaining functional. In contrast, the large abdominal muscle does not undergo atrophy during moulting. Crustacean muscles have two major fibre types that have previously been identified in *Cherax destructor* claw muscle; slow contracting fibres with long sarcomeres, and fast contracting fibres with short sarcomeres. All muscle fibre types are involved in moult-induced atrophy as well as moulting-associated fibre elongation, although different mechanisms appear to operate in each fibre type. In the present study, histochemical staining techniques have been developed to demonstrate contractile speed, fatigue resistance and fibre size of yabby muscle fibres. These techniques were used to determine the occurrence and distribution of fibre types and sub-types in the claw and abdomen during intermoult and to compare the fibre characteristics of moult-induced atrophic claw muscle with non-atrophic abdominal muscle over the moult cycle.

Respiratory responses of intertidal crab eggs to changes in oxygen tensions

N. Leelapiyanart¹ and H. H. Taylor²

¹Department of Aquatic Science, Burapha University, Chonburi, Thailand;

²Department of Zoology, University of Canterbury, Christchurch, New Zealand
nongnud@bucc4.buu.ac.th

In response to declining oxygen tension, pre-hatching eggs of *H. rotundifrons* and *C. lavauxi* exhibited characteristics of oxygen conformers and regulators respectively. The P_{crit} value for late stage eggs of *C. lavauxi* was approximately 6 kPa (45 mm Hg). The lactate concentration in eggs of *H. rotundifrons* increased from about 0.602 mmol.l⁻¹ in normoxic eggs to about 10.20 mmol.l⁻¹ after 3h hypoxic exposure. The rate of lactate accumulation during hypoxia suggests that the eggs are not sustaining metabolism by anaerobiosis, but are allowing metabolic depression. Pre-hatching eggs of *H. rotundifrons* and *C. lavauxi* exhibit no ventilatory or locomotory movements, and limited circulatory function. We hypothesize that the greater dependence of oxygen uptake on ambient PO_2 in the larger *H. rotundifrons* eggs reflects a greater diffusion limitation. This interpretation is supported by calculations of PO_2 gradients in a simplified model. This model showed that the centre of *Heterozius* eggs becomes anoxic at quite moderate external hypoxia, thus limiting total oxygen uptake. For *Cyclograpsus*, the centre does not become anoxic until severe external hypoxia prevails.

Acetylcholine controls surfactant secretion by type II pneumocytes isolated from Australian lungfish and American bullfrogs

Olga V. Lopatko¹, Philip G. Wood¹, Jean M. P. Joss², Allan W. Smits³ and Christopher B. Daniels¹.

¹Department of Physiology, University of Adelaide, Adelaide, SA 5000; ²School of Biological Sciences, Macquarie University, Sydney, NSW 2109; ³Quinnipiac College, Mt Carmel Avenue, Hamden, Connecticut, USA 06518-1940
pwood@physiol.adelaide.edu.au

The factors that control the release of pulmonary surfactant (PS) in non-mammalian vertebrates are largely unknown. In mammals, changes in ventilatory pattern and input from the sympathetic nervous system predominantly control surfactant secretion, while the parasympathetic nervous system has no effect. However, we have found that alterations in ventilatory pattern have no effect on surfactant release in lizards, but both the sympathetic and parasympathetic nervous systems stimulate surfactant phospholipid (PL) release. We examined the effects of the major autonomic neurotransmitters on PL release by type II pneumocytes isolated from lungs of the Australian lungfish, *Neoceratodus forsteri*, and the North American bullfrog, *Rana catesbiana*. Type II cells were isolated from each species by 'panning' whole lung cell suspensions over IgG coated bacteriologic plates. Cells were cultured for 15h in phosphate-free DMEM and PL secretion monitored in response to the application of adrenaline (Ad, 100 μ M) or acetylcholine (ACh, 100 μ M). Only ACh stimulated PL release from lungfish type II cells (an increase of 14%), but both Ad and ACh stimulated PL release from isolated bullfrog type II cells (Ad: 108% increase; ACh: 106% increase). Application of atropine blocked the ACh - elicited release of PL in both species. The Ad-stimulated release of PL from bullfrog pneumocytes was abolished by the presence of alprenolol. ACh also triggered the differential release of disaturated PL from lungfish and bullfrog type II cells. It appears that, *in vivo*, acetylcholine, and therefore the parasympathetic nervous system, may regulate the surfactant system in non-mammalian vertebrates, controlling surfactant release and composition. Adrenergic influences on the surfactant system may have developed subsequent to the radiation of the tetrapods.

Funded by the Australian Research Council and University of Adelaide and George Murray Overseas Travel Grants.


Development of the pulmonary surfactant system in a bird and reptile

Sandra Orgeig, Sonya D. Johnston, Olga V. Lopatko and Christopher B. Daniels

Department of Physiology, University of Adelaide, Adelaide, SA 5005
sjohnston@physiol.adelaide.edu.au

Little is known about the maturation of the pulmonary surfactant system in non-mammalian vertebrates. Unlike mammals, birds and reptiles hatch from eggs; often a lengthy and exhausting process whereby the animal commences lung clearance and pulmonary ventilation by "pipping" through the shell membranes. We examined the composition of pulmonary surfactant in the developing lungs of the chicken, *Gallus gallus*, and the bearded dragon, *Pogona vitticeps*. Lung tissue was collected from chicken embryos at day 14, 16, 18 (pre-pip) and 20 (post-pip) of incubation and from 1 day post-hatch, 3 week and adult animals. SPA mRNA was detected in lung tissue from chickens of all stages using Northern blot analysis. Chickens were lavaged at days 16, 18 (pre-pipped) and 20 (post-pipped) of incubation and 1 day post-hatch. Bearded dragons were lavaged at day 55 (pre-pipped), days 57-60 (post-pipped) and days 58-61 (post hatched). Total phospholipid (PL), disaturated phospholipid (DSP) and free cholesterol (CHOL) were measured in lavage fluid of both species. Total PL increased throughout incubation in both species (day 16 chick: 0.26 ± 0.09 μ g/mg dry lung weight (dlwt), n=4 to hatch: 0.93 ± 0.24 μ g/mg dlwt, n=9) and (day 55 bearded dragon: 0.39 ± 0.08 μ g/mg dlwt, n=4 to hatch: 4.42 ± 0.84 μ g/mg dlwt, n=7). DSP was not measurable at day 16 of incubation in the chick embryo. However the percentage of DSP/PL increased from day 18 (pre-pipped) to day 20 (post-pipped). As both PL and DSP increased with development, but CHOL remained unchanged during incubation in the chicken embryo, the CHOL to PL and CHOL to DSP ratios decreased in chick embryos respectively: (day 16: $39.53 \pm 6.72\%$, n=4 to hatch: $8.61 \pm 0.55\%$ n=9) and (day 18: $146.90 \pm 50.04\%$, n=8 to hatch $35.64 \pm 4.91\%$, n=9). This indicates that the CHOL and PL components are differentially regulated. The CHOL to PL ratio is higher in embryonic chickens than any other animal studied to date. The changes in lipid composition during development of the pulmonary surfactant system in the chicken and bearded dragon mimic those of the mammal despite the difference in birth strategy.

Supported by the Australian Research Council



Hyperlipaemic emu serum is clarified for radioimmunoassay by polyethyleneglycol (PEG) precipitation


Judith Van Cleeff, Margaret Blackberry and Graeme Martin
 Animal Science Department, University of Western Australia, Nedlands, WA 6907
 jvanclee@agric.uwa.edu.au

Emu serum collected during spring is often hyperlipaemic, and females laying in winter have high circulating concentrations of yolk proteins. The lipid or vitellogenin content in serum prevents accurate radioimmunoassay. Serum lipids are sequestered within lipoproteins, so we have adopted a method developed for analysis of lipoproteins to clarify samples. A solution of PEG (45% in water) is added to samples to precipitate large molecular weight proteins by steric exclusion. Centrifugation separates clarified serum from the lipoprotein precipitate. PEG treatment clarified 97% of emu serum samples (n=500) collected from May to October. The nature of the precipitate varied according to sex and season. We identified four types of precipitate: 1) thin, cohesive (48%); 2) floating, fatty (16%; 67% male); 3) large volume, loose (26%; 96% from breeding-season females); 4) floating fatty layer and thin precipitate (5%). Samples which were not clarified (4%) were from egg-laying females. Of type 2 precipitates, 71% were found in October (non-breeding season). Residuals after PEG precipitation of emu serum spiked with tracer-conjugated hormones were 87, 89, 93, 100, and 100% for insulin, LH, glucagon, prolactin, and testosterone, respectively. Hyperlipaemic samples being prepared for radioimmunoassay can be clarified with minimal hormone loss using PEG precipitation.

Diet and assimilation in the robber crab, *Birgus latro* (L.)

Joanne Wilde and Peter Greenaway
 School of Biological Sciences, UNSW, Sydney, NSW 2052
 joanne@unsw.edu.au

From observations of the natural diet on Christmas Island, it was clear that *Birgus latro* principally ate fruits and seeds rich in storage carbohydrates (*Pandanus*, *Ochrosia ackeringae*) and lipids (*Aleurites moluccana*, *Calophyllum inophyllum*, *Arenga listeri*) but also predated on other crabs and scavenged animal carcasses. The capacity to assimilate the major components of these diets was assessed in the laboratory using 3 artificial diets rich in plant polysaccharides, plant lipids and crab tissue respectively. Assimilation of dry matter, carbohydrate and lipids were high on all diets. Assimilation efficiencies (AE's) of dry matter were similar for the three diets ($71.7 \pm 0.7\%$, $75.5 \pm 1.3\%$ and $64.7 \pm 5.1\%$, for carbohydrate, fat and crab diets respectively), but AE's of lipid were significantly different ($87.4 \pm 1.7\%$, $96.03 \pm 0.7\%$ and $70.8 \pm 3.7\%$, respectively). Near complete assimilation of polysaccharides occurred in the carbohydrate and fat diets ($98.9 \pm 0.2\%$ and $98.06 \pm 0.7\%$, respectively) but assimilation was significantly lower in the crab diet ($89.4 \pm 4.5\%$). Dry matter intake was affected inversely by the N concentration of the diets and as lowest ($0.86 \pm 0.08 \text{ g.kg}^{-1}.\text{day}^{-1}$) on the crab diet ($5.4 \text{ mmol N.g}^{-1}$), intermediate ($99 \pm 0.61 \text{ g.kg}^{-1}.\text{day}^{-1}$) on the fat diet ($1.2 \text{ mmol N.g}^{-1}$) and highest ($4.31 \pm 0.52 \text{ g.kg}^{-1}.\text{day}^{-1}$) on the carbohydrate diet ($0.9 \text{ mmol N.g}^{-1}$). Fibre was a relatively small component of the carbohydrate and fat diets but assimilation of hemicellulose ($49.5 \pm 0.7\%$ and $64.8 \pm 2.9\%$, respectively) and cellulose ($16.1 \pm 4.3\%$ and $53.9 \pm 3.5\%$, respectively) was quite high. The rate of passage, measured with particulate tracers, was slow with a mean retention time of $27.3 \pm 7.22 \text{ h}$ but as defecation is frequent this value may be overestimated.



Registrants

<i>Surname</i>	<i>First names</i>	<i>Address</i>	<i>email</i>	<i>phone</i>	<i>fax</i>
Adams	Dominic	Institute of Natural Resources (Ecology), Massey University, Palmerston North, New Zealand	D.C.Adams@massey.ac.nz	64 06 350 5315	350 5623
Andersen	Niels	Division of Anatomy & Physiology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001	niels.andersen@utas.edu.au	03 6226 2655	6226 2679
Anderson	Trevor	Department of Aquaculture, James Cook University, Townsville, QLD 4811	Trevor.Anderson@jcu.edu.au	07 4781 5586	4781 4585
Baldwin	John	Department of Biological Sciences, Monash University, Clayton, VIC 3168	john.baldwin@sci.monash.edu.au	03 9905 5613	9905 5613
Beal	Michel	School of Biological Science, University of New South Wales, Sydney, NSW 2052	m.beal@unsw.edu.au	02 9385 2116	9385 1558
Bech	Claus	Department of Zoology, Norwegian University of Science and Technology, N-7034 Trondheim Norway	claus.bech@chemblo.ntnu.no		
Bothge	Philip	Department of Anatomy & Physiology, University of Tasmania, GPO Box 252-24, Hobart, TAS 7001	p_bothge@utas.edu.au	03 6226 2687	6226 2679
Blache	Dominique	Animal Science, University of Western Australia, Nedlands, WA 6907	dblache@agnc.uwa.edu.au	08 9380 3587	9380 1040
Bradshaw	Don	Department of Zoology, University of Western Australia, Nedlands WA 6907	Don.Bradshaw@uwa.edu.au	08 9380 3769	9380 1463
Bradshaw	Felicity	Department of Zoology University of Western Australia, Nedlands WA 6907	fjb@cyllene.uwa.edu.au	08 9380 2248	9380 1029
Buttmer	Bill	Department of Biological Sciences University of Wollongong, Wollongong, NSW 2522	buttmer@uow.edu.au	02 4221 4459	4221 4135
Celi	Pietro	Faculty of Agriculture (Animal Science) The University of Western Australia, Nedlands, WA 6907	pceli@agnc.uwa.edu.au	08 9380 7026	9380 1040
Chappell	Mark	Department of Biology, University of California at Riverside, Riverside, CA 92521 USA	chappell@citrus.ucr.edu		
Cutler	Lanelle	School of Natural Sciences, Edith Cowan University, 2 Bradford Street, Mt Lawley WA 6050	lanellec@hotmail.com	08 9370 6138	9370 6103
Donald	John	Department of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217	jdonald@deakin.edu.au	03 5227 2097	5227 2022

<i>Surname</i>	<i>First names</i>	<i>Address</i>	<i>email</i>	<i>phone</i>	<i>fax</i>
Edwards	Ashley	School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001	Ashley.Edwards@utas.edu.au	03 6226 2592	6226 2745
Edwards	Peta	School of Biological Science, University of New South Wales, Sydney, NSW 2052	02 9385 3262	93821 558	
Elias	John	Department of Biological Sciences, Monash University, Clayton, VIC 3168	john.elias@sci.monash.edu.au	03 9375 4209 9905 5758	
Geiser	Fritz	Zoology, School of Biological Sciences, University of New England, Armidale, NSW 2351	fgeiser@metz.une.edu.au	02 6773 2868	6773 3814
Gemmell	Robert	Department of Anatomical Sciences, University of Queensland, St. Lucia, QLD 4072	R.Gemmell@mailbox.uq.edu.au	07 3365 2626	3365 1299
Greenaway	Peter	School of Biological Sciences, University of New South Wales, Sydney, NSW 2052	p.greenaway@unsw.edu.au	02 93852067	9385 1558
Guppy	Michael	Department of Biochemistry, University of Western Australia, Nedlands, WA 6907	mguppy@cylene.uwa.edu.au	08 9380 3331	9380 1148
Hoh	Joe	Department of Physiology, University of Sydney, Sydney, NSW 2006	joeh@physiol.usyd.edu.au	02 9351 4267	9351 2058
Hope	Perdita	Department of Medicine, Level 6 Services Building, Royal Adelaide Hospital, Adelaide, SA 5000	phope@medicine.adelaide.edu.au	08 8222 4329	8222 3870
Hulbert	Tony	Biological Sciences, University of Wollongong, Wollongong, NSW 2522	a.hulbert@uow.edu.au	02 4221 3437	4221 4135
Hume	Ian	Biological Sciences A08, University of Sydney, Sydney, NSW 2006	ianhume@bio.usyd.edu.au	02 9351 2369	9351 4119
Johnston	Sonya	Department of Physiology, University of Adelaide, Adelaide, SA 5005	sjohnston@physiol.adelaide.edu.au	08 8303 4157	8303 3356
Jones	Sue	School of Zoology, University of Tasmania, GPO Box 252-05, Hobart, TAS 7001	S.M.Jones@utas.edu.au	03 6226 2592	6226 2745
Joss	Jean	School of Biological Sciences, Macquarie University, Sydney, NSW 2109	jjoss@ma.bio.mq.edu.au	02 9850 8137	9850 8245
Lamey	Tina	School of Natural Sciences, Edith Cowan University, 2 Bradford Street, Mt Lawley, WA 6050	tinalamey@hotmail.com	08 9370 6138	9370 6103
Leeapiyanart	Nongnud	Department of Aquatic Science, Burapha University, Chonburi, Thailand	nongnud@bucc4.buu.ac.th	66 038 745 900 xtn 3023	390 354
Malecki	Irek	Faculty of Agriculture (Animal Science), University of Western Australia, Nedlands, WA 6907	imalecki@agnc.uwa.edu.au	08 9380 7025	9380 1040
McAllan	Bronwyn	Human Biology, School of Biological Sciences, University of New England, Armidale, NSW 2351	bmcallan@metz.une.edu.au	02 6773 2048	6773 3267



<i>Surname</i>	<i>First names</i>	<i>Address</i>	<i>email</i>	<i>phone</i>	<i>fax</i>
Munns	Suzanne	Department of Zoology University of Melbourne Parkville VIC 3052	s.munns@pgrad.unimelb.edu.au	03 9344 6244	9344 7909
Newbound	Dennysa	Zoology Department The University of Western Australia Nedlands 6907, WA 6907	dnewb@cyllene.uwa.edu.au	08 9380 3970	9380 1029
Nicol	Stewart	Division of Anatomy & Physiology, University of Tasmania, GPO Box 252-24, Hobart, TAS 7001	s.c.nicol@utas.edu	03 6226 2655	6226 2679
O'Shea	Jamie	Department of Zoology, University of Western Australia, Nedlands, WA 6907	jeoshea@cyllene.uwa.edu.au	08 9380 2242	9380 1029
Pakay	Julian	Department of Biochemistry, University of Western Australia, Nedlands, WA 6907	jlpakay@cyllene.uwa.edu.au	08 9380 2242	9380 1148
Patak	Annette	School of Natural Sciences, Edith Cowan University, 2 Bradford Street, Joondalup, WA 6050	a.patak@cowan.edu.au	08 9370 6658	9370 6103
Paterson	Brian	Centre for Food Technology, Queensland Department of Primary Industry, 19 Hercules Street, Hamilton, QLD 4007	patersonb@dpi.qld.gov.au	07 3406 8577	3406 8698
Pearson	James	Department of Electrical and Electronic Engineering, Muroran Institute of Technology, Muroran, Mizumoto-cho 27-1, 050-8585 JAPAN	jpearson@muroran-it.ac.jp	81 143 465 533	465 501
Riono	Agung	Faculty of Agriculture (Animal Science) University of Western Australia, Nedlands, WA 6907	ariono@cyllene.uwa.edu.au	08 9380 2976	9380 1040
Roberts	Alan	Department of Zoology, University of Western Australia, Nedlands, WA 6907	aroberts@cyllene.uwa.edu.au	08 9380 3962	9380 1029
Roberts	Julie	Animal Physiology, School of Rural Science and Natural Resources, University of New England, Armidale, NSW 2351	jroberts@metz.une.edu.au	02 6773 2506	6773 3234
Semia	Conrad	Dept Physiology & Pharmacology, University of Queensland, St Lucia, QLD 4072	C.Semia@mailbox.uq.edu.au	07 3365 3180	3365 8228
Spanoghe	Patrick	School of Biomedical Sciences, Curtin University of Technology, GPO Box 1987, Perth, WA 6001	rspanogh@cc.curtin.edu.au	08 9266 4132	9266 2342
Taylor	Harry	Department of Zoology, University of Canterbury, Private Bag 4800, Christchurch New Zealand	h.taylor@zool.canterbury.ac.nz	64 3 364 2861	364 2024
Thompson	Anton	Department of Zoology, University Western Australia, Nedlands, WA 6907	apt@cyllene.uwa.edu.au	08 9380 1477	9380 1029



<i>Surname</i>	<i>First names</i>	<i>Address</i>	<i>email</i>	<i>phone</i>	<i>fax</i>
Thompson	Graham	Centre for Ecosystem Management, Edith Cowan University, 100 Joondalup Drive, Joondalup, WA 6027	G.Thompson@cowan.edu.au	08 9400 5427	9400 5840
Tomanovic Toop	Rada Tes	16 Andell Place, Redcliffe, WA 6104 School of Biological and Chemical Sciences, Deakin University, Geelong, VIC 3217	08 9277 5567 ttoop@deakin.edu.au	02 5227 1260	5227 2022
van Aardt	Willie	Zoology Department, Potchefstroom Universiteit, Potchefstroom, Private Bag X6001, South Africa 2520	drkja@puknet.puk.ac.za	27 18 299 2562	299 2370
van Cleeff	Judith	Department of Animal Science, University of Western Australia, Nedlands, WA 6907	jvanclee@agnc.uwa.edu.au	08 9380 2976	9380 1040
Veitch	Colleen	Department of Anatomical Sciences, University of Queensland, St Lucia, QLD 4072	C.Veitch@mailbox.uq.edu.au	07 3365 2961	3365 1299
Wilde	Joanne	School of Biological Sciences, University of New South Wales, Sydney, NSW 2052	joanne@unsw.edu.au	02 93852011	9385 1558
Withers	Philip	Department of Zoology, University of Western Australia, Nedlands WA 6907	pwithers@cyllene.uwa.edu.au	08 9380 2235	9380 1029
Wittert	Gary	Department of Medicine, Level 6 Services and Teaching Building, Royal Adelaide Hospital, Adelaide, SA 5000	gwittert@medicine.adelaide.edu.au	08 8222 5502	82233870
Wood	Philip	Department of Physiology, University of Adelaide, Adelaide, SA 5000	pwood@physiol.adelaide.edu.au	08 8303 4157	8303 3356



Weekdays

Route No.		Cottesloe	Claremont	Crawley	Busport I
72	am	5:45	5:59	6:06	6:12
72		6:20	6:34	6:41	6:47
71		6:40	6:54	7:01	7:07
71		-	-	-	7:27
71		7:03	7:18	7:26	7:37
72		-	-	-	7:47
70		7:15	7:31	7:39	7:52G
72		-	-	-	7:57
70		7:25	7:41	7:49	8:02G
72		-	-	-	8:07
70		7:35	7:51	7:59	8:12G
72		-	-	-	8:17
70		7:47	8:03	8:11	8:24G
71		7:59	8:15	8:23	8:36
72		8:11	8:28	8:36	8:49
71		8:33	8:48	8:56	9:09
71		8:53	9:08	9:16	9:29
73		-	-	-	9:40
71		9:25	9:40	9:48	9:59
73		-	-	-	10:20
71		9:55	10:10	10:18	10:29
71		10:25	10:40	10:48	10:59
73		-	-	-	11:00
72		10:55	11:10	11:18	11:29
73		-	-	-	11:40
71		11:25	11:40	11:48	11:59
73		-	-	-	12:20
71	pm	11:55	12:10	12:18	12:29
71		12:25	12:40	12:48	12:59
73		-	-	-	1:10
71		12:55	1:10	1:18	1:29
72		1:25	1:40	1:48	1:59
73		-	-	-	2:15
71		1:55	2:10	2:18	2:29
72		2:24	2:40	2:48	2:59
73		-	-	-	3:12
71		2:54	3:10	3:18	3:29
73		-	-	-	3:42
72		3:19	3:35	3:43	3:54
72		3:29B	3:45	3:53	4:04
71		-	-	4:08J	4:19
73		-	-	-	4:22
71		3:59	4:15	4:23	4:34
71		-	-	4:38J	4:49
73		-	-	-	4:52
72		4:30	4:45	4:53	5:04
71		4:40	4:55	5:03	5:14
73		-	-	-	5:22
72		4:52	5:07	5:15	5:26
72		5:10	5:25	5:33	5:44
73		-	-	-	5:57
72		5:40	5:55	6:03	6:12
72		5:55	6:10	6:18	6:27
72		6:15	6:30	6:38	6:47
72		6:50	7:04	7:11	7:17
72		7:20	7:34	7:41	7:47
72		8:00	8:14	8:21	8:27
72		-	-	9:06J	9:12
72		9:58	10:12	10:19	10:25
72		10:35	10:49	10:56	11:02
72		11:13	11:27	11:34	11:40


Saturdays

Route No.		Cottesloe	Claremont	Crawley	Busport
72	am	7:05	7:19	7:26	7:32
72		7:41	7:55	8:02	8:08
72		8:10	8:24	8:31	8:37
72		8:40	8:55	9:03	9:12
73		9:15	9:30	9:38	9:47
72		9:50	10:05	10:13	10:22
73		10:25	10:40	10:48	10:57
72		11:00	11:15	11:23	11:32
73	pm	11:35	11:50	11:58	12:07
72		12:10	12:25	12:33	12:42
73		12:45	1:00	1:08	1:17
72		1:20	1:35	1:43	1:52
73		1:55	2:10	2:18	2:27
72		2:30	2:45	2:53	3:02
73		3:05	3:20	3:28	3:37
72		3:40	3:55	4:03	4:12
73		4:15	4:30	4:38	4:47
72		4:50	5:05	5:13	5:22
72		5:25	5:40	5:48	5:57
72		6:00	6:15	6:23	6:32
72		7:10	7:24	7:31	7:37
72		8:15	8:29	8:36	8:42
72		9:40	9:54	10:01	10:07
72		10:55	11:09	11:16	11:22

Sundays and Public Holidays







Route No.		Cottesloe	Claremont	Crawley	Busport
72	am	9:25	9:39	9:46	10:00
72		11:05	11:19	11:26	11:32
72		11:50	12:04	12:11	12:17
72	pm	1:25	1:39	1:46	2:00
72		2:35	2:49	2:56	3:02
72		4:25	4:39	4:46	5:00
72		5:05	5:19	5:26	5:32
72		6:55	7:09	7:16	7:30



Map routes 70, 71, 72, 73, 74, 75, 76, 77



Map routes 200, 201, 202, 208, 209, 556

Legend

-  Train Station with Train Connection
-  Train Station
-  Special Event Train Station
-  Bus Station
-  Bus Route Terminus
-  Timed Stop



Bus from Claremont						
Route No.	Claremont	Nedlands	West Perth	Perth	Domestic Airport	
209	7:00A	7:10A	7:20A	7:36A	-	
202	7:15	7:33	7:42	7:47	8:28	
201	7:20A	7:36A	-	7:53A	-	
209	7:30	7:43	7:55	8:00	8:41	
202	7:40A	7:58A	8:07A	8:19A	-	
201	7:45	8:01	-	8:17	9:02	
209	7:55A	8:08A	8:20A	8:38A	-	
202	8:05B	8:23B	8:32B	8:37B	9:19	
201	8:15A	8:31A	-	8:53A	-	
202	8:30B	8:45B	8:56B	9:02B	9:42	
208	8:55B	9:10B	9:25B	9:30B	10:12	
201	9:25B	9:41B	-	9:57B	10:42	
202	10:00BC	10:15BC	10:26BC	10:32BC	11:18	
208	10:25B	10:40B	10:55B	11:00B	11:42	
201	11:02BD	11:18BD	-	11:34BD	12:23	
202	11:30B	11:45B	11:56B	12:02B	12:42	
208	11:59BC	12:14BC	12:29BC	12:34BC	1:20	
201	pm 12:30B	12:46B	-	1:02B	1:47	
202	1:00BD	1:15BD	1:26BD	1:32BD	2:18	
208	1:25B	1:40B	1:55B	2:00B	2:42	
201	2:02BC	2:18BC	-	2:34BC	3:23	
202	2:20E	2:35E	2:50	2:56BD	3:42	
200	-	-	-	3:20B	4:04	
201	3:02BC	3:18BC	-	3:34BC	4:25	
202	3:18E	3:33E	3:48	3:54	4:33	
208	3:35C	3:50C	4:05C	4:12C	4:57	
202	3:49D	4:04D	4:15D	4:21D	5:06	
209	3:55A	4:08A	4:20A	4:36A	-	
202	4:12	4:27	4:38	4:44	5:25	
201	4:26C	4:42C	-	4:58C	5:46	
209	4:39	4:52	5:04	5:09	5:50	
202	4:46D	5:01D	5:12D	5:18D	6:03	
201	5:00	5:16	-	5:32	6:14	
209	5:19	5:32	5:44	5:49	6:28	
202	5:30	5:48	5:57	6:02	6:41	
202	6:08	6:23	6:31	6:35	7:10	
200	6:40	6:55	7:05	7:09	7:44	
200	7:25	7:40	7:50	7:54	8:29	
200	8:50	9:05	9:15	9:19	9:54	
200	10:10	10:25	10:35	10:39	11:14	





Saturdays



Bus from Claremont						
202	am	7:00	7:15	7:23	7:27	8:02
201		7:40B	7:54B	-	8:08B	8:47
208		8:15B	8:27B	8:40B	8:44B	9:18
202		9:00B	9:15B	9:23B	9:27B	10:05
201		9:45B	9:59B	-	10:13B	10:52
208		10:20B	10:32B	10:45B	10:49B	11:23
202		11:00B	11:15B	11:23B	11:27B	12:05
201		11:45B	11:59B	-	12:13B	12:52
208	pm	12:25B	12:37B	12:50B	12:54B	1:28
202		1:10B	1:25B	1:33B	1:37B	2:15
200		2:00B	2:15B	2:25B	2:29B	3:07
200		3:05B	3:20B	3:30B	3:34B	4:12
200		4:00B	4:15B	4:25B	4:29B	5:07
200		4:50B	5:05B	5:15B	5:19B	5:57
200		6:05	6:20	6:30	6:34	7:09
200		7:30	7:45	7:55	7:59	8:34
200		9:55	10:10	10:20	10:24	10:59
200		11:01	11:16	11:26	11:30	12:05

Sundays and Public Holidays

Bus from Claremont						
200	am	9:14	9:29	9:39	10:00	10:35
200		11:14	11:29	11:39	12:00	12:35
200	pm	1:14	1:29	1:39	2:00	2:35
200		2:49	3:04	3:14	3:18	3:53
200		4:14	4:29	4:39	5:00	5:35
200		-	-	-	6:02	6:37
200		6:44	6:59	7:09	7:30	8:05

Map routes 200, 201, 202, 208, 209, 556

- Legend**
-  Train Station with Train Connection
 -  Train Station
 -  Special Event Train Station
 -  Bus Station
 -  Bus Route Terminus
 -  Timed Stop



Exciting changes at *Physiological Zoology* in 1999 and beyond

Physiological Zoology
changes its title in 1999 to
Physiological and Biochemical Zoology
and introduces its online edition

Since its inception in 1928, *Physiological Zoology* has been devoted to the study and publication of significant and timely research in circulation, metabolic energetics and regulation, nutrition and digestion, temperature adaptation, sensory physiology, acid-base balance and neural coordination, and other fields.

• **Title Change in 1999**

The University of Chicago Press is pleased to announce that beginning with Volume 72 (January/February 1999) *Physiological Zoology* will be published under its new title, *Physiological and Biochemical Zoology*. This title change will not alter the focus or editorial direction of the journal, but rather, convey more accurately *PZ*'s traditional emphasis of original research in environmental, adaptational, and comparative physiology and biochemistry. Editing and production standards will continue at the same high standards to which you are accustomed.

• **Available Online in 1999—**

<http://www.journals.uchicago.edu/PBZ>

In 1999 a new era begins as *Physiological and Biochemical Zoology* launches its electronic edition—accessible via the world-wide web at <http://www.journals.uchicago.edu/PBZ>. Using the online version of *PBZ* will be an easy way to access the latest investigations and developments, timely research and methods, and important theories and commentaries as soon as they are published. Subscribers will continue to receive the bimonthly paper edition as well as the electronic edition, which is updated weekly. The electronic edition is included with subscriptions to the paper edition at no additional charge. Immediate access and powerful search features make the electronic edition of *PBZ* an invaluable tool.

• **Beyond 1999**

For more than seventy years of publication, *Physiological Zoology* has served researchers, faculty, teachers, and students alike. As *Physiological and Biochemical Zoology* enters the next millennium, the journal will continue building upon the traditions and relationships that *PZ* has developed throughout its long and prestigious history.

Physiological Zoology—exploring all aspects of comparative physiology and biochemistry

Edited by Gregory K. Snyder, *University of Colorado at Boulder*

Published by The University of Chicago Press for the
Division of Comparative Physiology and Biochemistry of the Society for Integrative and
Comparative Biology

One-year subscription rates (includes electronic access): Institution \$328.00; Individual \$69.00; Student (attach copy of valid ID) \$41.00; SICB Individual Member \$55.00. Outside USA, please add \$11.00 postage. Contact The University of Chicago Press for airmail rates. Canadian residents, please also add 7% GST. Visa and MasterCard payments accepted. Credit card customers may fax their orders to (773) 753-0811. To order, send check or credit card information (acct #, exp. date, telephone number, and signature) to:

The University of Chicago Press
Journals Division, Dept. SF9SA, P.O. Box 37005, Chicago, IL 60637 USA
For more information, please visit our website at <http://www.journals.uchicago.edu/PZ/>

Announcing exciting changes at *Physiological Zoology*

Physiological Zoology changes its title in 1999 to
Physiological and Biochemical Zoology
and introduces its online edition

• **Title Change in 1999**

The University of Chicago Press is pleased to announce that beginning with Volume 72 (January/February 1999) *Physiological Zoology* will be published under its new title, *Physiological and Biochemical Zoology*. This title change will not alter the focus or editorial direction of the journal, but rather, convey more accurately PZ's traditional emphasis of original research in environmental, adaptational, and comparative physiology and biochemistry.

• **Available Online in 1999—<http://www.journals.uchicago.edu/PBZ>**

In 1999 a new era begins as *Physiological and Biochemical Zoology* launches its electronic edition—accessible via the world-wide web at <http://www.journals.uchicago.edu/PBZ>. Using the online version of PBZ will be an easy way to access the latest investigations and developments, timely research and methods, and important theories and commentaries as soon as they are published. Subscribers will continue to receive the bimonthly paper edition as well as the electronic edition, which is updated weekly. The electronic edition is included with subscriptions to the paper edition at no additional charge. Immediate access and powerful search features make the electronic edition of PBZ an invaluable tool.

• **Beyond 1999**

As *Physiological and Biochemical Zoology* enters the next millennium, the journal will continue building upon the traditions and relationships that PZ has developed throughout its long and prestigious history.

Physiological Zoology—exploring all aspects of comparative physiology and biochemistry

Edited by Gregory K. Snyder, University of Colorado at Boulder

Published by The University of Chicago Press for the

Division of Comparative Physiology and Biochemistry of the Society for Integrative and Comparative Biology

One-year subscription rates (includes electronic access): Institution \$328.00; Individual \$69.00; Student (attach copy of valid ID) \$41.00; SICB Individual Member \$55.00. Outside USA, please add \$11.00 postage. Contact The University of Chicago Press for airmail rates. Canadian residents, please also add 7% GST. Visa and MasterCard payments accepted. Credit card customers may fax their orders to (773) 753-0811. To order, send check or credit card information (acct #, exp. date, telephone number, and signature) to:

The University of Chicago Press

Journals Division, Dept. SF9SA, P.O. Box 37005, Chicago, IL 60637 USA.

For more information, please visit our website at <http://www.journals.uchicago.edu/PZ/>

10/98

CBP COMPARATIVE BIOCHEMISTRY & PHYSIOLOGY

AN INTERNATIONAL JOURNAL - PARTS A, B, C

An Official Journal of the

**Australian & New Zealand
Society for Comparative
Physiology & Biochemistry**

Japanese Society for
Comparative physiology
& Biochemistry

Society for Experimental
Biology

Canadian Society of
Zoologists (CPB Section)

European Society for
Comparative Physiology
& Biochemistry

Society for Experimental
Biology

South American Society
for Comparative Physiology
& Biochemistry

Society for Integrative &
Comparative Biology
(Formerly the American
Society of Zoologists)

Editors-in-Chief: **Peter W. Hochachka &
Thomas P. Mommsen**

CBP Editorial Office, University of British Columbia,
6371 Crescent Road, Vancouver BC, Canada V6T 1Z2
Fax: 604 228-0408

Review Editor: **Pat J. Walsh**

NIEHS Marine & Freshwater Biomedical Science Center,
Rosentiel School of Marine & Atmospheric Science,
University of Miami, 4600 Rickenbacker Causeway, Miami,
FL 33149-4617, USA
Fax: 305 361-4600

AIMS & SCOPE:

The journal publishes original articles emphasizing comparative and environmental aspects of the physiology, biochemistry, molecular biology, pharmacology, toxicology and endocrinology of animals. Adaptation and evolution as organizing principles are encouraged. Studies on other organisms are considered if approached in a comparative text.

Part A. Molecular and Integrative Physiology. This section deals with molecular, cellular, integrative, and ecological physiology. Topics include bioenergetics, circulation, development, excretion, ion regulation, neurobiology, nutrition, respiration, and thermal biology. Study on regulatory mechanisms at any level of organization such as signal transduction and cellular interactions and control of behaviour are encouraged.

Part B. Biochemistry and Molecular Biology. This section covers biochemical and molecular biological aspects of metabolism, enzymology, regulation, nutrition, signal transduction, promoters, gene structure and regulation, metabolite and cell constituents, macromolecular structures, adaptational mechanisms and evolutionary principles.

Part C. Pharmacology, Toxicology, and Endocrinology. This section is concerned with chemical and drug action at different levels of organization, biotransformation of xenobiotics, mechanisms of toxicity, including reactive oxygen species and carcinogenesis, hormones and receptors, natural products chemistry, and signal transduction. A molecular approach to these fields is encouraged.

Members of ANZCPB may
subscribe at the following
rates:

Part A, ISSN: 0300-9629
Volumes 122-124, 1999
12 issues per annum
US\$ 118

Part B, ISSN: 0305-0491
Volumes 122-124, 1999
12 issues per annum
US\$ 118

Part C, ISSN: 0742-8413
Volumes 122-124, 1999
9 issues per annum
US\$ 93

Parts A, B & C
Volumes 122-124, 1999
33 issues per annum
US\$248

Prices include postage & are subject to
change without notice

Special
Reduced Rate
to ANZSCP
Members!

INDEXED/ABSTRACTED IN: CABS

(Current Awareness in Biological Sciences),
*Curr. Cont. ASCA, Aqua Abstr., Curr. Cont./Life Sci.,
Curr. Cont. ISI/BIOMED Database, Reference Update,
Curr. Cont. SCISEARCH Data*



FROM ELSEVIER SCIENCE...
SCIENCE PUBLISHER TO THE WORLD



Calling all professionals interested in Biotechnology!
Get FREE access to ContentsDirect, the on-line, e-mail service for Biotechnology-related journals

If you have e-mail, you can have access to tables of contents listings for the following Elsevier Science journals! Approximately two to four weeks prior to the publication of each issue, **ContentsDirect** provides you with the contents listing for that issue - *directly to your PC*.

You get:


- journal title
- volume and issue number
- issue publication date
- article title
- author names
- page numbers

If you would like to be added to this fast, free and direct service, visit the **ContentsDirect** web site.

That's all it takes!

No cost!
No obligation!
•
Guaranteed pre-publication - Delivered 2-4 weeks prior to publication
•
Easy to use - Just open your e-mail

- Advanced Drug Delivery Reviews
- Biomedical Materials
- Biosensors and Bioelectronics
- Biotechnology Advances
- Drug Discovery Today
- Enzyme and Microbial Technology
- European Journal of Pharmaceutics and Biopharmaceutics
- Genetic Analysis: Biomolecular Engineering
- International Journal of Biodeterioration and Biodegradation
- International Journal of Pharmaceutics
- Journal of Molecular Catalysis B: Enzymatic
- Journal of Fermentation and Bioengineering
- Journal of Biotechnology
- Mutation Research: Fundamental and Molecular Mechanisms of Mutagenesis
- Mutation Research Genomics
- Pharmaceutical Science & Technology Today
- Process Biochemistry
- Trends in Biotechnology
- Trends in Biochemical Sciences



ContentsDirect

Register now!

ContentsDirect

www.elsevier.nl/locate/ContentsDirect

NB if you can't access the web - please note that you can still register by sending an E-mail to cdsubs@elsevier.co.uk

**Get quality research in fermentation
and bioengineering from the journal
devoted to its study**

Journal of Fermentation and Bioengineering

Published by the Society of
Fermentation and Bioengineering,
Japan. Distributed outside Japan by
Elsevier Science

Chairman: T. Kobayashi

International Editors:

T. Imanaka, (*Editor-in-Chief, Japan*)

Osaka University, Osaka, Japan, Jochem Büchs,
(*Germany, Europe*) **Aachen University of
Technology, Aachen, Germany, P.L. Rogers**,
(*Australia, Oceania*) **The University of New South
Wales, Australia and H.Y. Wang**, (*USA, America*)
The University of Michigan, Ann Arbor, MI, USA

Devoted to the advancement and
dissemination of knowledge concerning
fermentation technology, biochemical
engineering, food technology and
microbiology.

Journal of Fermentation and Bioengineering is
indexed/abstracted in:
BIOSIS, Chemical Abstracts, Current Contents.



ContentsDirect Now Available.

Elsevier's FREE ContentsDirect service is now available
for this journal.

This alerting service sends the table of contents of each
issue to your PC 2-4 weeks prior to publication.

To register as a new user, visit the ContentsDirect Web site
at <http://www.elsevier.com/locate/contentsdirect>

1998 Subscription Information

Volumes 85-86 (12 issues)

ISSN 0922-338X

Subscription price: NLG 1713.00 /US\$984.00

Visit our website for more details about the journal:
<http://www.elsevier.nl/locate/jfermbio>

For more information or to place an order, contact

For customers in the Americas and Canada
Elsevier Science
P.O. Box 945, New York
N.Y. 10159-0945, U.S.A.
Tel: (+1) 212 633 3730
[Toll Free number for
North-American customers:
1-888-4ES-INFO (437 4636)]
Fax: (+1) 212 633 3680
E-mail: usinfo-f@elsevier.com

For customers in all other locations
Elsevier Science
P.O. Box 211
1000 AE Amsterdam
The Netherlands
Tel: (+31) 20 485 3757
Fax: (+31) 20 485 3432
E-mail: nlinfo-f@elsevier.nl




LINK




Experimental Biology Online - EBO

Online journal


<http://link.springer.de/journals/ebo/>



Ensure that you don't miss your access to EBO in 1998 and enter your subscription now!



Now also available:
The archive edition of the complete contents of the electronic journal "Experimental Biology Online - EBO" from the years 1996 and 1997



ISSN: 1430-3418

Experimental Biology Online - EBO is a new, multimedia electronic journal sponsored by three international biological societies (Society for Experimental Biology/SEB, Society for Integrative and Comparative Biology/SICB and the Australian and New Zealand Society for Comparative Physiology and Biochemistry/ANZSCPB) with a membership of over 6,000 worldwide. It provides a powerful medium for rapid, refereed short research papers in the general field of experimental biology, ranging from the molecular and cellular boundaries in cell and plant biology to the integrative study of whole organisms including the experimental analysis of behaviour. **EBO** specializes in short, topical research papers in experimental biology organized in three sections.

Features at a glance

- All papers are submitted and refereed by peer-review electronically either via e-mail or anonymous FTP server. This usage of the Internet at every step of the reviewing and publication process leads to exceedingly short publication times of a few weeks only from submission to appearance
- The journal is accessible on the World Wide Web, with long-term access guaranteed
- Full color figures plus additional material, e.g. picture, sound or movie files are also included.
- Language of publication: English
- Electronic format: HTML and multimedia supplements
- **EBO** is accessible on the World Wide Web via Springer's information service **LINK** (<http://link.springer.de>).

Subscription prices in 1998

- Institutional subscription:** DM 198,- / US \$ 122.00
- Personal subscription:** DM 98,-
- Members of affiliated societies:** DM 78,-

Editors-in-Chief

- Editor-in-Chief Animal Biology Section**
Prof. Christopher R. Bridges
University of Dusseldorf, FRG
e-mail: bridges@uni-duesseldorf.de
- Editor-in-Chief Plant Section**
Prof. Dale Sanders
University of York, UK
e-mail: ds10@york.ac.uk
- Editor-in-Chief Cell Biology Section**
Prof. Adam Curtis
University of Glasgow, UK
e-mail: a.s.g.curtis@bio.gla.ac.uk

C.R. Bridges/D. Sanders/A. Curtis

EBO - Experimental Biology Online Annual (vol. 1996/1997)

1998. Annual with CD-ROM for PC.
ISBN 3-540-64146-7
DM 198,-
Special price for members of the SEB and affiliated societies: DM 98,-

System requirements for archive CD-ROM:

IBM compatible PC with 80486 SX processor, 8 MB RAM, VGA (640 x 480) video card set to 256 colours, mouse, double-speed CD-ROM player or faster, Windows 95

...and now take a closer look online at EBO and...

Search it - See it! - Hear it! - Try it!

<http://link.springer.de/journals/ebo/>

- Animal Biology
- Plant Biology
- Cell Biology

SEB
SOCIETY FOR EXPERIMENTAL BIOLOGY



Springer
electronic media



From the contents - recently published papers...

Second-messenger-induced signalling events in pollen tubes of Papaver rhoeas (Plant Biology Section), B.K. Dröbak, F.C.H. Franklin, P.J. Shaw, G.M. Calder, A.J. Trewavas, A.C. Allan and V.E. Franklin-Tong

An improved method for screening Frankia viability in strain Ccl3 (Plant Biology Section), N.R. McEwan and C.T. Wheeler

Experimental simulation in behavioral ecology: a multimedia approach with the spatial searching simulation RattleSnake (Animal Biology Section), D. Duvall, D. Chiszar, R.A. Mintzer, and N.O. Roster

Effects of Diazinon on bluegill sunfish, Lepomis macrochirus, gills: scanning electron microscope observations (Cell Biology Section), H. M. Dutta, J. S. D. Munshi, P. K. Roy, N. K. Singh, L. Moiz and S. Adhikari

The multiple saluble malate dehydrogenase of Hoplias malabaricus (Erythrinidae, Characiformes) (Animal Biology Section), M.R. Aquino-Silva, M.L.B. Schwantes and A.R. Schwantes

Quantitative imaging of intact cells and tissues by multi-dimensional confocal fluorescence microscopy (Cell Biology Section), M.D. Fricker, C.-M. Chow, R.J. Errington, M. May, J. Mellor, A.J. Meyer, M. Tlalka, D.J. Vaux, J. Wood and N.S. White

Effect of fasting and thermal acclimation on metabolism of juvenile axolotls (Ambystoma mexicanum) (Animal Biology Section), Louis N. Irwin, Karen A. Talentino, and Denise A. Caruso

Cell behaviour in tubes (Cell Biology Section), Mathis Riehle, Diana Ferris, Douglas Hamilton and Adam Curtis

Editors-in-Chief

(Animal Biology Section) Christopher R. Bridges, University of Düsseldorf/Germany, bridges@uni-duesseldorf.de

(Plant Biology Section) Dale Sanders, University of York/UK, dsio@york.ac.uk

(Cell Biology Section) Adam Curtis, University of Glasgow/UK, a.s.g.curtis@bio.gla.ac.uk

Associate Editors

SICB (ASZ): Joan D. Ferraris, NIH, Bethesda/USA, jdf@helix.nih.gov

ANZSCP: Stewart Nicol, University of Tasmania/Australia, s.c.nicol@physiol.utas.edu.au

Editorial Board Animal Biology Section

John Altringham, j.d.altringham@leeds.ac.uk

Pat Butler, p.j.butler@bham.ac.uk

Steven Cork, s.cork@dwe.csiro.au

Geoffrey M. Coast, g.coast@biol.bbk.ac.uk

Andrew Cossins, cossins@pop1.liv.ac.uk

Joachim W. Deitmer, deitmer@rhrk.uni-k.de

David Duvall, duvalld@okway.okstate.edu

David H. Evans, dhesfish@zoo.ufl.edu

Gerd Flik, gertflik@sci.kun.nl

Robert Full, rjfull@garnet.berkeley.edu

Gerd Gaede, ggaede@botzoo.uct.ac.za

James W. Hicks, jhicks@uci.edu

Peter Lutz, lutz@acc.fau.edu

Kathy Packard, kpackard@huliy.colostate.edu

Steve Perry, sperry@oreo.uottawa.ca

John Fleng Steffensen, mbljfs@inet.uni-c.dk

Steve Morris, stevems@bio.usyd.edu.au

Kenneth Storey, kbstorey@ecs.carleton.ca

Harry Taylor, h.taylor@zool.canterbury.ac.nz

Andre Toulmond, toulmond@sb-roscoff.fr

Michele G. Wheatly, mwheatly@desire.wright.edu

Roddy Williamson, rwil@pml.ac.uk

Editorial Board Plant Biology Section

David Baulcombe, david.baulcombe@bbsrc.ac.uk

Dan Cosgrove, dcosgrove@psu.edu

Jeff Dangl, dangl@email.unc.edu

Bill Davies, w.davies@lancaster.ac.uk

Simon Gilroy, sxg12@email.psu.edu

Wilhelm Gruissem, gruisssem@nature.berkeley.edu

Roger Hangarter, rhangart@bio.indiana.edu

Dirk Inze, diinz@gengenp.rug.ac.be

Bill Lucas, wjlucas@aggie.ucdavis.edu

Helen Ougham, helen.ougham@bbsrc.ac.uk

Steve Tyerman, steve.tyerman@flinders.edu.au

Editorial Board Cell Biology Section

J. Bereiter-Hahn, bereiter-hahn@zoology.uni-frankfurt.d400.de

Patrick Aebischer, patrick.aebischer@chuv.hospvd.ch

P. Bongrand, bongrand@infobiogen.fr

John Bryant, j.a.bryant@exeter.ac.uk

Julian Dow, j.a.t.dow@bio.gla.ac.uk

Graham Dunn, gad@helios.rai.kcl.ac.uk

Clive Evans, c.evans@auckland.ac.nz

Dennis Francis, fraancid@cardiff.ac.uk

Albert Harris, akharris@med.unc.edu

Jeff Hubbell, hubbell@cheme.caaltech.edu

Evamaria Kinne, evamaria.kinne@mpi-dortmund.mpg.de

Eisuke Nishida, enishida@virus.kyoto-u.ac.jp

Jack Steele, jenny@pelican.dbe.csiro.au

Vickery Trinkaus-Randall, vickery@med-biochem.bu.edu

ORDER FORM

Please bill me

Please order through your bookseller or from:

Empty box for ordering details

Tel.: ++49 (0) 30-8 27 87-358 · Fax: ++49 (0) 30-8 27 87-448
E-mail: subscriptions@springer.de · http://www.springer.de
Springer-Verlag, Postfach 31 13 40, D-10643 Berlin

Yes, please enter my subscription* for EBO, starting with 1998
ISSN 1430-3418 Title No. 898:

- Institutional subscription: DM 198,- / US \$ 122.00
 Personal subscription: DM 98,-
 Members of affiliated societies: DM 78,- (SEB, SICB, ANZSCP)

Please send me:

EBO - Experimental Biology Online Annual (vol. 1996/1997) (ISBN 3-540-64146-7)

- DM 198,- DM 98,- (special price for members of affiliated societies)

LINK-No. (if available)

e-mail address

Name

Street

City/ZIP-Code

Country

Date/Signature

* Please note: I realize that I may cancel this subscription within ten days by writing to the address on the order form, whereby the postmark date will suffice as proof that the cancellation was made within the deadline. I confirm my understanding of this by my second signature.

Second Signature:

39302/3.98.d&p.WM Prices are subject to change without notice.
Customers in EU: Customers, please state your VAT-identification-Number if applicable.

http://link.springer.de/journals/ebo/



Access & Licensing

A wide range of material is available on the LINK server free of charge, such as table of contents and abstracts. Access to complete articles is restricted to print subscribers. The procedure is somewhat different for institutional subscribers, individual subscribers and societies but in all cases there is a single access and identification point for all journals of the Springer Group of Companies.

LINK Licences:

Institutional subscribers can choose between:

Basic LINK Licence: Access charge included in print to all subscribed LINK journals subscription in 1998 and 1999 for three simultaneous users.

Unlimited LINK Licence per subscribed journal (prices upon request).

Institutional users are identified by user-ID, password and the IP addresses of their computers. Where an IP address covers exactly those computers whose users have access to the printed editions of the subscribed journals, we also offer access control by "IP only".

Please complete onscreen the contract at:

<http://link.springer.de/orders/rest.htm> and send it via fax or mail to Springer-Verlag.

Personal LINK Licences

give one named individual access to her/his subscribed journals. The user is identified by user-ID and password. Access is possible from any computer - from work, home or when travelling. As an individual user with a personal subscription please use the online registration form at: http://link.springer.de/access_ind/index.htm

Tutorial

LINK Tutorial always provides the new information on how to get access, whom to contact, frequently asked questions, technical details etc.

This service is designed to help you through LINK. Please have a look at: <http://link.springer.de/tutorial/>

Equipment

There are no special hardware requirements to get access to LINK: Just a computer - either PC, Mac or UNIX-based system - and a connection to the Internet.

That might be a connection via your company or university network or a dial-up connection by modem or ISDN.

For playing audio files, a sound system (e.g. Soundblaster) has to be installed on your computer.

Nevertheless, some programs for viewing multimedia supplements are only available for Windows 95.

LINK Orders

Our helpdesk provides support by phone, fax and e-mail.

Operating hours are 9am to 5pm MET.

Orders from USA, Canada, Mexico

Online ordering
<http://link.springer-ny.com/order-na.htm>


Technical and customer support
LINK Helpdesk
Springer-Verlag New York Inc.
175 Fifth Avenue
New York, NY 10010, U.S.A.
Tel. +1/212/460-1612
FAX +1/212/473 6272
e-mail: link@springer-ny.com

Orders from Other Countries

Online ordering:
<http://link.springer.de/orders/rest.htm>

Technical and customer support
LINK Helpdesk
Springer-Verlag
Tiergartenstr. 17
D-69121 Heidelberg
Tel. +49/180 5656665
FAX +49/6221/487-120
e-mail: helpdesk@link.springer.de

The Visionary
Information Service



<http://link.springer.de>
link.springer-ny.com

E Electronic Journals

E Electronic Versions of Printed Journals and Books

E P Electronic Supplementary Material

<http://link.springer.de>
link.springer-ny.com



LINK - The Visionary Information Service

The LINK information service offers electronic versions of most printed journals and an increasing number of books from the Springer Group of Companies on the Internet, thus linking printed and electronic products. LINK is bringing a wide variety of information right to your desktop, whether you are a scientist, lecturer, librarian, information broker or student.

LINK's content is always expanding. You will find almost all journals - both with purely electronic and online versions of print journals - and multimedia supplements. LINK contains an increasing number of electronic books, moderated discussion forums, information on meetings and conference schedules. LINK is fully supported - around the world - by dedicated staff who can answer questions and troubleshoot any problem.

Online Libraries


LINK creates an extensive digital marketplace called "The Forum for Science", which is organised into fully searchable Online Libraries as shown below. These Online Libraries

allow direct access to the table of contents, abstracts and full text of the journals published by Springer-Verlag and soon by other publishing houses.

Ten Online Libraries are the core of LINK:

- Chemical Sciences Online Library
- Computer Science Online Library
- Economics Online Library
- Engineering Online Library
- Environmental Sciences Online Library
- Geosciences Online Library
- Life Sciences Online Library
- Mathematics Online Library
- Medicine Online Library
- Physics & Astronomy Online Library

The entirety of Online Libraries is called "Forum for Science"



LINK Alert

Subscribers and nonsubscribers of our journals will automatically via e-mail and free of charge receive the table of contents with direct links to the respective abstracts as soon as a new issue of a journal you are interested in is available in LINK. You can either select from a list of all the journals and other products and developments that might interest you or you can click on the LINK Online Libraries and then choose your area of interest. <http://link.springer.de/alert/>

Online First

With Online First articles can be published in electronic form weeks before distribution of the print journal - even before the issue and page numbers have been assigned. This is not a preprint service, the publications are in their final form. Online First articles are citeable by the "Digital Object Identifier", or DOI, a unique and persistent identification code, included in both the print and electronic versions.

LINK Search

LINK offers three different search possibilities, each with Boolean Operators. Searches are carried out in either one selected library or the whole data set.

LINK Search: Search in bibliographic records, abstracts of the articles and the general information about the journals.

LINK Power Search: Structured search in specified bibliographic fields; journal title, title, author, affiliation, abstract, keyword, publication year

LINK Global Search: Search in all data available in the LINK Service - including full articles. The result of a search request is a list of hits, containing all articles that match the search conditions. From the list you get a direct link to the abstract page.


Both Power Search and Global Search offer additional search terms as BUT NOT, NEAR and FOLLOWED BY and ranking

according to: relevance, number of occurrences of the search term, presence in a specific document region.


Moderated Forum

The introduction of moderated forums will allow scientific discussions with the editors or authors, and are excellent tools for the systematic incorporation of user feedback. This means a forum for discussing new ideas, controversial or even contentious papers, and the numerous, often diverging opinions, terms, and regulations which are in existence or for which proposals are in preparation. Such a forum for public discussion aims at gradually bringing clarity and therefore contributing to international understanding and meaningful measures on an intercontinental level.

A journal's outfit



The index page of each journal leads to the tables of contents and information about the journal




All abstracts are displayed in HTML format

Most of the articles are available in different file formats. The most widely used format is the Portable Document Format (.pdf). In pdf-format the articles on your monitor look like printed ones, thus this file format is ideal for hardcopies. You can access and print the papers directly from your desktop. Other formats for delivery of full articles are Postscript, TeX and HTML. The format type differs from journal to journal, according to the contents. For exam-

ple: Mathematical formulas might not be displayed correctly using HTML, so TeX is the format of choice in this case. In LINK we offer access to supplementary material such as: Color Images, Sound, Video, and Data sets. You will find the necessary plug-ins (e.g. Adobe Acrobat Reader, Quicktime-Player, IBM-technexplorer) free of charge on our LINK server at: <http://link.springer.de/help/helpers.htm>

Most of the articles are available in different file formats



The Royal Society of Western Australia has a World Wide Web (WWW) home page that is maintained by the Honorary Editor on a server courtesy of Dr Graham Thompson and Edith Cowan University. The URL address of the home page is <http://leisuresci.fhhs.ac.cowan.edu.au/RSWA/>. The web pages were written and are maintained using Microsoft FrontPage 98® and are best viewed with Microsoft Internet Explorer 4® at 1024x840.

The home page of the RSWA provides information concerning:

- Council Members
- Meetings information
- Proceedings [updated monthly]
- Journal Information, including history, instructions to authors, tables of content, and publications for sale;
- Symposia
- History of the Society
- Medallists: 1924-1997
- Library Holdings
- Membership Application Form
- Constitution
- Links to other sites of interest – including other Royal Societies

The RSWA web pages link to three searchable databases with information on Publications of the Royal Society, its Library holdings, and its Councils. A simple search-engine allows wild-card searching of specific fields of the databases and returns all records found, as a simplified subset of the database.

- **PUBLICATIONS:** This database summarises the publications of the Royal Society of Western Australia and its precursor Societies (over 2250 records). It contains all papers published as well as other information for each issue, including information from the title page, the inside front cover, and the inside and outside of back cover, as well as additional pages such as Proceedings, Annual Society Report, Treasurer's Report, Table of Contents, Index, Corrigenda and Addenda, etc. Fields included in specific searches are Author and Year, and Key-word is searched for in all fields. The fields returned for the records found are: Year, Authors, Title, Journal, Reference.
- **LIBRARY HOLDINGS:** This database includes all journal and book listings of the holdings of the Library of the Royal Society of Western Australia. It only comprises a single field, the index number and full citation of the holding.
- **COUNCIL:** This database details the membership of Society's Council, including the precursor Societies to the Royal Society of Western Australia (over 95 records). It contains information for all Society Councils (President, Vice presidents, Secretaries, Treasurer, Editor and Assistants, Librarian and Assistants, Journal Manager, general members), year of office, Patron, Vice Patron, Auditors, and Other Appointments. All fields are included in the search and returned.

Search by Author.

Fifth International Congress of Comparative Physiology and Biochemistry (ICCPB99)

CPB99

Calgary, Alberta, Canada
August 23-28, 1999

SPECIAL PLENARY LECTURE
Dr. Michael Smith

PLENARY SPEAKERS

Dr. R.E. Peter (Canada)
Dr. P.W. Hochachka (Canada)
Dr. M. Konishi (Japan/USA)
Dr. Y. LeMaho (France)
Dr. M. McFall-Ngai (USA)

THEMES

A. Biotechnology and Biochemical Control
B. Cell and Molecular Control
C. Endocrinology and Development
D. Physiology and Regulatory Control
E. Neurosciences

SCIENTIFIC OBJECTIVE

The primary objective of the Congress is to bring together comparative physiologists and biochemists worldwide to promote debate and discussion on topics of wide current interest. Throughout we will provide a forum for interdisciplinary participation and cross fertilization of ideas. A satellite symposium on paleophysiology is also planned.

SYMPOSIA AND WORKSHOPS (Details available at Website)

A. BIOCHEMICAL CONTROL

- Antifreeze proteins: evolution, regulation and mechanisms of action
- Oxygen binding and related proteins: structure, function and evolution
- Toxin versus energy source: physiological and biochemical strategies of adaptation to sulfide-rich environments
- Osmoregulatory plasticity: regulators and conformers
- Research and development in NO
- Molecular and mechanical design of structural materials
- Protein Structural Motifs

B. CELL AND MOLECULAR CONTROL

- Molecular cell biology of osmoregulation
- Molecular control of muscle plasticity - opportunities for comparative biology

Web Site:
<http://www.acs.ucalgary.ca/~iccpb99>

HONORARY PRESIDENT
Dr. W.S. Hoar

ADDRESS

The Secretariat, ICCPB99
Conference Management Services
Olympic Volunteer Centre
1833 Crowchild Trail N.W., Calgary, Alberta, Canada T2M 4S7
Fax: (403) 284-1184 E-Mail: iccpb99@acs.ucalgary.ca

SPONSORS

Canadian Society of Zoologists
International Union of the Biological Sciences

PARTICIPATING SOCIETIES

Canadian Society of Zoologists
European Society for Comparative Physiology & Biochemistry
Society for Integrative Comparative Biology
Japanese Society for Comparative Physiology & Biochemistry
Society for Experimental Biology
Societe de Physiologistes
Deutsche Zoologische Gesellschaft
Australian and New Zealand Society for Comparative Physiology and Biochemistry
American Physiological Society Section for Comparative Physiology and Biochemistry
Russian Physiological Society Section for Comparative Biochemistry

- Gene regulation by environmental stress
- Environmental contamination and cell signalling
- Molecular responses to thermal stress
- Hibernation and daily torpor as the periodic expression of genes
- Cell culture in the physiology of lower vertebrates
- Drosophila as a model system for comparative physiologists and biochemists

C. ENDOCRINOLOGY AND DEVELOPMENT

- Comparative aspects of hormone receptors and signal transduction
- Environmental endocrine disruptors and their impact on growth and development
- Novel control mechanisms in vertebrate neuroendocrine systems
- Integrated control of brain-pituitary-peripheral axes
- Novel aspects on the biochemistry and physiology of lipoproteins in insects
- Evolution of viviparity: the physiological issues, recent research and

- Integrative crustacean biology: evolutionary trends
- The eye lens: function, structure, development and evolution
- Reproductive technologies: a comparative approach


D. PHYSIOLOGY AND REGULATORY CONTROL

- Water, land and air: Unifying principles in locomotion: Water
- Water, land and air: Unifying principles in locomotion: Land
- Ontogeny, diversity and adaptation of respiratory control
- Specialized circulatory structures in fish: Secondary circulation and pseudobranch
- Cardiac and Vascular strategies of lower vertebrates
- Cardiovascular reflexes and their neural pathways in invertebrates
- Physiological constraints on the ecology and behavior of marine predators
- Foraging in evolution: Optimal feeding and digestion in animals
- Nitrogen metabolism in fish: Molecular and evolutionary aspects of

- Physiological thermoregulation in anurans
- Seasonal effects on physiological function
- Physiological challenges of aquacultured animals
- New optical methods in physiology and biochemistry: Imaging from gene to organism

E. NEUROSCIENCES

- Physiological effects of cardioactive neuropeptides in invertebrates
- Myoactive peptides in invertebrate animals: structures, evolution and function
- Insect brain: Simple design of sophisticated behaviour
- How animals in nature use their sense of smell
- Diversity of visual pigments
- Molecular aspects of circadian clock systems
- Evidence and possible roles for signalling proteins in extrasomal neuronal compartments



Foraging Lake, Banff National Park

Photo courtesy of Robert Berlin